

TRI-BASIC

Containing Inert Diluent

DIRECTIONS FOR USE: Tri-basic Bordeaux Powder should be used in accordance with recommendations of State or Federal Authorities for Basic Copper Sulfate compounds containing the equivalent of 12.75% metallic copper. The following general directions are supplied by us gratuitously and are believed to be reliable but are in no way guaranteed.

For Use On	Control These Diseases or Pests	Amount of Diluent Required per Gallon	Amount of Diluent Required When
FRUITS			
Small Cherries	Brown Rot Leaf spot	4 tablespoonsful plus 1/2 tablespoonful of Hydrated Spray Lime per gallon of water.	Start to apply as soon as leaves appear and repeat every 10 days.
Grapes	Black Rot Downy mildew	1/2 tablespoonful with 1/2 tablespoonful of Hydrated Spray Lime per gallon of water.	Start to apply at end of dormant season every 10 days.
Apples	Bitter Rot Blotch Scab	1/2 tablespoonful with 1/2 tablespoonful of Hydrated Spray Lime per gallon of water.	After leaves open repeat every 10 days.
Gooseberries Currants	Anthraxnose Leaf Spot	4 1/2 tablespoonful with 1/2 tablespoonful of Hydrated Spray Lime per gallon of water.	After leaves open repeat every 10 days.
VEGETABLES			
Potatoes	Early blight Late blight	1/2 tablespoonful per gallon of water.	Apply when plants are 16 inches high. Repeat every 10 to 14 days.
Tomatoes	Anthraxnose Early blight Late blight	1/2 tablespoonful per gallon of water.	When plants first set out repeat every 10 days.
Cantaloupes Cucumbers Squash Watermelons	Angular leaf spot and scab Anthraxnose Downy mildew	1/2 tablespoonful per gallon of water.	As soon as plants begin to leaf repeat every 10 days.
Greens	Early blight Late blight	1/2 tablespoonful per gallon of water.	Apply in seed bed every 10 days.
TREES			
Apple Oak Maple Birch	Anthraxnose Leaf spot Bark beetle	1/2 tablespoonful per gallon of water.	Spray as soon as leaves appear. Repeat every 10 days.

CAUTION: Harmful if Swallowed

ANTIDOTE: In case of inhalation or swallowing give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Follow with milk or whites of eggs beaten with water. In any case call a physician immediately.

Tri-basic Bordeaux Powder is not recommended for early application on varieties of apples susceptible to copper russetting.

Store in a dry cool area, completely out of reach of children or pets. Do not contaminate any body of water.

Do not reuse empty container. Destroy it by burying with waste and burning. Stay away from smoke or fumes.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
No. 305

Containing Inert Diluent

for use in the orchard and garden

ACTIVE INGREDIENT
Copper in Basic Copper
Sulfate (expressed
as Basic)
12.75%
INERT INGREDIENTS
87.25%
TOTAL 100.00%

CAUTION
KEEP OUT OF THE
REACH OF CHILDREN
See side panel for other cautions.



EPA REG. NO. 5887-41-AA
NET WEIGHT 1 LB.


CEP

3-30-73

THE FEDERAL
SIDE AND BOX
CONTAINING PERS
NO. 305

5887-41

043073



**for use in the orchard
and garden**



DIRECTIONS FOR USE: Tri-basic Bordeaux Powder should be used in accordance with recommendations of State or Federal Authorities for Basic Copper Sulfate compounds containing the equivalent of 12.75% metallic copper. The following general directions are supplied by us gratuitously and are believed to be reliable but are in no way guaranteed.

FRUITS			
Sour Cherries	Brown Rot Leaf spot	4 tablespoonsful plus 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	Start to apply as soon as leaves ap- pear and repeat ev- ery 10 days.
Grapes	Black Rot Downey mildew	7 tablespoonsful with 4 tablespoonsful of Hydrated Spray Lime per gallon of water.	Start to apply at end of dormant sea- son every 10 to 14 days.
Apples	Bitter Rot Blotch Scab	3 tablespoonsful with 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	After leaves open re- peat every 7 to 10 days.
Gooseberries Currants	Anthrachnose Leaf Spot	4 1/2 tablespoonsful with 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	After leaves open cover leaves thor- oughly.
VEGETABLES			
Potatoes	Early blight Late blight	14 tablespoons per gallon of water.	Apply when plants are 4-6 inches high. Repeat every 10 to 14 days.
Tomatoes	Anthrachnose Early blight Late blight	14 tablespoons per gallon of water.	When plants first set out, repeat ev- ery 10-14 days.
Cantaloupes Cucumbers Squash Watermelons	Angular leaf spot and scab Anthrachnose Downey mildew	7 1/2 tablespoonsful per gallon of water.	As soon as plants begin to vine, re- peat every 7 days.
Celery	Early blight Late blight	11 1/4 tablespoons per gallon of water.	Apply in seed bed every 7-10 days.
TREES			
Linden Oak Maple Elm	Anthrachnose Leaf spot Elm leaf curl	6 to 7 tablespoons per gallon of water.	Spray as leaves un- curl. Repeat every 2-3 weeks.

Depending on use, this package will make up to 25 gallons of spray.

CAUTION: Harmful if Swallowed

ANTIDOTE: In case of inhalation or swallowing give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Follow with milk or whites of eggs beaten with water. In any case call a physician immediately.

Tri-basic Bordeaux Powder is not recommended for early application on varieties of apples susceptible to copper russetting.

Store in a dry cool area, completely out of reach of children or pets. Do not contaminate any body of water.

Do not reuse empty container. Destroy it by burying with waste and burning. Stay away from smoke or fumes.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
OO. 303

0-73

5887-41

Black
Leaf

Bordeaux Powder

for use in the orchard
and garden

A FUNGICIDE for use on grapes,
sour cherries, cantaloupes, squash,
apples, gooseberries, potatoes,
tomatoes, cucumbers, celery,
watermelons.

Bordeaux Powder

DIRECTIONS FOR USE: Tri-basic Bordeaux Powder should be used in accordance with recommendations of State or Federal Authorities for Basic Copper Sulfate compounds containing the equivalent of 12.75% metallic copper. The following general directions are supplied by us gratuitously and are believed to be reliable but are in no way guaranteed.

FRUITS			
Sour Cherries	Brown Rot Leaf spot	4 tablespoonful plus 2 tablespoonful of Hydrated Spray Lime per gallon of water.	Start to apply as soon as leaves appear and repeat every 10 days.
Grapes	Black Rot Downy mildew	7 tablespoonful with 4 tablespoonful of Hydrated Spray Lime per gallon of water.	Start to apply at end of dormant season every 10 to 15 days.
Apples	Bitter Rot Blotch Scab	3 tablespoonful with 2 tablespoonful of Hydrated Spray Lime per gallon of water.	After leaves open repeat every 7 to 10 days.
Gooseberries Currants	Anthraxnose Leaf Spot	4 1/2 tablespoonful with 2 tablespoonful of Hydrated Spray Lime per gallon of water.	After leaves open cover leaves thoroughly.
VEGETABLES			
Potatoes	Early blight Late blight	14 tablespoons per gallon of water.	Apply when plants are 4-6 inches high. Repeat every 10 to 14 days.
Tomatoes	Anthraxnose Early blight Late blight	14 tablespoons per gallon of water.	When plants first set out, repeat every 10-14 days.
Cantaloupes Cucumbers Squash Watermelons	Angular leaf spot and scab Anthraxnose Downy mildew	7-9 tablespoonful per gallon of water.	As soon as plants begin to vine, repeat every 7 days.
Celery	Early blight Late blight	11-14 tablespoons per gallon of water.	Apply in seed bed every 7-10 days.
TREES			
Linden Oak Maple Elm	Anthraxnose Leaf spot Elm leaf curl	6 to 7 tablespoons per gallon of water.	Spray as leaves uncurl. Repeat every 2-3 weeks.

Depending on use, this package will make up to 25 gallons of spray.

CAUTION: Harmful if Swallowed

ANTIDOTE: In case of inhalation or swallowing give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Follow with milk or whites of eggs beaten with water. In any case call a physician immediately.

Tri-basic Bordeaux Powder is not recommended for early application on varieties of apples susceptible to copper russetting.

Store in a dry cool area, completely out of reach of children or pets. Do not contaminate any body of water.

Do not reuse empty container. Destroy it by burying with waste and burning. Stay away from smoke or fumes.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.


DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
00-303

TEIX

0-73

INSECTICIDE

5887-41



for use in the orchard
and garden

A Fungicide. Use on grapes,
sour cherries, cantaloupes, squash,
apples, gooseberries, peaches,
tomatoes, cucumbers,
watermelons.

ACTIVE INGREDIENT	
Copper (in Basic Copper	
Sulfate) Expressed	12.75%
as metallic	87.25%
INERT INGREDIENTS	
TOTAL	100.00%



CAUTION
KEEP OUT OF THE
REACH OF CHILDREN
See side panel for other cautions.

EPA REG. NO. 5887-41-AA
NET WEIGHT 1 LB.

Bordeaux Powder

DIRECTIONS FOR USE: Tri-basic Bordeaux Powder should be in accordance with recommendations of State or Federal Authorities for Basic Copper Sulfate compounds containing the equivalent of 12.75% metallic copper. The following general directions are supplied by us gratuitously and are believed to be reliable but are in no way guaranteed.

FRUITS			
Small Cherries	Brown Rot Leaf spot	4 tablespoonsful plus 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	Start to apply as soon as leaves appear and repeat every 10 days.
Grapes	Black Rot Downy mildew	7 tablespoonsful with 4 tablespoonsful of Hydrated Spray Lime per gallon of water.	Start to apply at end of dormant season every 10 to 15 days.
Apples	Bitter Rot Blotch Scab	3 tablespoonsful with 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	After leaves open to post every 7 to 10 days.
Gooseberries Currants	Anthraxnose Leaf Spot	4 tablespoonsful with 2 tablespoonsful of Hydrated Spray Lime per gallon of water.	After leaves open cover leaves thoroughly.
VEGETABLES			
Potatoes	Early blight Late blight	14 tablespoons per gallon of water.	Apply when plants are 4-6 inches high. Repeat every 10 to 14 days.
Tomatoes	Anthraxnose Early blight Late blight	14 tablespoons per gallon of water.	When plants first set out, repeat every 10-14 days.
Cantaloupes Cucumbers Squash Watermelons	Angular leaf spot and scab Anthraxnose Downy mildew	7.5 tablespoonsful per gallon of water.	As soon as plants begin to vine, repeat every 7 days.
Celery	Early blight Late blight	11-14 tablespoons per gallon of water.	Apply in seed bed every 7-10 days.
TREES			
Linden Oak Maple Elm	Anthraxnose Leaf spot Elm leaf curl	6 to 7 tablespoons per gallon of water.	Spray as leaves uncurl. Repeat every 2-3 weeks.

Depending on use, this package will make up to 25 gallons of spray.

CAUTION: Harmful if Swallowed

ANTIDOTE: In case of inhalation or swallowing give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Follow with milk or whites of eggs beaten with water. In any case call a physician immediately.

Tri-basic Bordeaux Powder is not recommended for early application on varieties of apples susceptible to copper russetting.

Store in a dry cool area, completely out of reach of children or pets. Do not contaminate any body of water.

Do not reuse empty container. Destroy it by burying with waste and burning. Stay away from smoke or fumes.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
00-303

TEIX

0-73

5887-41