



Yellow - Applicant Copy

2/2

NOTIFICATION

SEP 1 1 2002



Tri-Basic Bordeaux Powder

- Controls Blights, Leaf Spot, Black Rot, Downy Mildew, Bud Rot and Other Listed Diseases
- Controls Palm Bud Rot
- Use on Listed Fruits, Vegetables, Flowers, Trees and Ornamentals



ACTIVE INGREDIENT:
Copper (in Basic Copper Salts) 12.75%
Expressed as Metallic Copper
INERT INGREDIENTS: 87.25%
TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for Additional Precautions.

NET CONTENTS 1 LB (454 g)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply as a spray as directed below.

USE PRECAUTIONS

REENTRY: Do not allow people or pets to enter treated area until sprays have dried.
Bordeaux Powder Fungicide is not compatible with dinitro compounds, dithiocarbamates, or lime sulfur. Do not apply during cool damp weather as "burning" or "scorching" of treated plants may occur. Young tender leaves of peach, plum, apple and rose are susceptible to spray injury. BORDEAUX POWDER FUNGICIDE is not recommended for early application on varieties of apples susceptible to copper russeting.

	DISEASE OR FUNGUS	HOW MUCH	WHEN
Hydrangea	Leaf Spot, Blight	6-9 tablespoons per gallon of water.	Apply as growth starts in Spring. Repeat as needed.
Willow, Dahlias, Tulips, Gladioli	Bacterial Blight	6-9 tablespoons per gallon of water.	Apply as growth starts in Spring. Repeat at 10 day intervals if plants have blight.
Hollyhock, Peonies	Anthracnose, Leaf Spot	6-9 tablespoons per gallon of water.	Spray after periods of wet weather.
Rose	Downy Mildew	6-9 tablespoons per gallon of water.	Spray every 2 weeks during growing season.
Banana	Black Leaf Spot	6-9 tablespoons per gallon of water.	Apply when new leaves appear. Repeat 2-3 times, 10 days apart.
Barberry	Canker	6-9 tablespoons per gallon of water.	Spray twice in Spring on twigs in Summer.
Isabel, Lilac	Leaf Blight	6-9 tablespoons per gallon of water.	Spray when new leaves appear. Repeat twice, 10 days apart.
Pines	Needle Blight	6-9 tablespoons per gallon of water.	Spray new growth in Spring.
Yew	Twig Blight	6-9 tablespoons per gallon of water.	Start when new growth in Spring. Repeat twice, one week apart.
Willow	Twig Blight, Black Canker	6-9 tablespoons per gallon of water.	Spray when new leaves are 1/4". Repeat twice, two weeks apart. Prune heavily to remove all second parts.
Abutilon, Cypripedium, Juncus, Delphinium	Leaf Blotch, Twig Blight, Cystenium Blight, Leaf Spot	6-9 tablespoons per gallon of water.	Apply in early Spring. Repeat at 2 week intervals as needed.
Blackberry, Aralia	Dieback	6-9 tablespoons per gallon of water.	Spray as new leaves appear. Repeat 2 weeks later. Prune and destroy affected twigs.
Rose	Palm Bud Rot	6-9 tablespoons per gallon of water.	Apply early in the Spring and every 2 weeks following periods of wet periods of wet weather as needed.
Scab Cherries	Brown Rot, Leaf Spot	4 tablespoons with 2 tablespoons of hydrated spray lime per gallon of water.	Start to apply as soon as leaves appear and repeat every 10 days.
Grape	Black Rot, Downy Mildew	7 tablespoons with 2 tablespoons of hydrated spray lime per gallon of water.	Start to apply at end of dormant season and repeat every 10-14 days.
Apple	Bitter Rot, Black, Scab	3 tablespoons with 2 tablespoons of hydrated spray lime per gallon of water.	Start to apply after leaves open and repeat every 7-10 days.
Gooseberries, Currants	Anthracnose, Leaf Spot	4 1/2 tablespoons with 2 tablespoons of hydrated spray lime per gallon of water.	Start to apply after leaves open. Cover leaves thoroughly. Repeat every 10-14 days.
Peaches	Leaf Curl, Brown Rot, Peach Blight, Shot Hole	4 tablespoons per gallon of water.	First spray for Brown Rot and Peach Blight should be applied before bud swell and in full pink bud stage. Do not apply after petal bud or after leaves are in bud as injury may occur.
Note: To control Leaf Curl application must be made before the foliage bud swells. As a dormant spray for Leaf Curl and Peach Blight, apply between Nov. 15 and Dec. 15 before rains begin. Over-spraying peaches may cause injury to tender foliage.			
Potatoes	Early Blight, Late Blight	14 tablespoons per gallon of water.	Apply when plants are 4-6 inches high. Repeat every 10-14 days.
Tomatoes	Anthracnose, Early Blight, Late Blight	14 tablespoons per gallon of water.	Apply when plants are first set out and repeat every 10-14 days.
Cantaloupes, Cucumbers, Squash, Watermelons	Angular Leaf Spot and Scab, Anthracnose, Downy Mildew	7-9 tablespoons per gallon of water.	Apply as soon as plants begin to vine and repeat every 7 days.
Celery	Early Blight, Late Blight	11-14 tablespoons per gallon of water.	Apply in seed bed every 7-10 days.
Indigo, Oak, Maple, Elm	Anthracnose, Leaf Spot, Elm Leaf Curl	6-7 tablespoons per gallon of water.	Spray as leaves unfold. Repeat every 2-3 weeks.

Depending on use, this package will make up to 25 gallons of spray.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes or clothing. May cause skin sensitization in certain individuals. Wash thoroughly after handling, before eating or using tobacco. Avoid contamination of food and feedstuffs. Food utensils such as measuring cups should not be used for food purposes after use in measuring this product.

FIRST AID			
If Inhaled:	• Move person to fresh air.	If In Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 min.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.		• Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.
If on Skin or Clothing:	• Call a poison control center or doctor for further treatment advice.		• Call a poison control center or doctor for treatment advice.
	• Take off contaminated clothing.	If Swallowed:	• Call a poison control center or doctor immediately for treatment advice.
	• Rinse skin immediately with plenty of water for 15-20 min.		• Have person sip a glass of water if able to swallow.
	• Call a poison control center or doctor for further treatment advice.		• Do not induce vomiting unless told to do so by the poison control center or doctor.
			• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

STORAGE & DISPOSAL

Store to be stored in the original container placed in areas inaccessible to children. **Pesticide Disposal:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Do not reuse this container. Place in trash or other for recycling if available.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

Buyer assumes responsibility for safety and use not in accordance with label directions.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-838-7378.

Distributed by
Valley Garden Supply, LLC
PO Box 585, St. Joseph, MO 64502-0585
(862) 864-6477
www.valleygardensupply.com

EPA Reg. No. 5887-41
EPA Est. No. 44616-J0-1
FMA#13004



NOTIFICATION

SEP 1 1 2002