

NET CONTENTS
ONE POUND



ACCEPTED

12-14-66

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC POISON PREVENTION
ED UNDER NO. 5887-43

CAUTION:

HARMFUL IF INHALED, SWALLOWED,
OR ABSORBED THROUGH THE SKIN.

Do Not Breathe Dust. Do Not Take In-
ternally. Avoid Prolonged or Repeated
Breathing of Spray.

Skin Contact With Any Insecticide May
be Harmful. Avoid Unnecessary Skin
Contact. Wash Hands Before Eating.

Note for Physician: Carbaryl is a moderate,
reversible, Cholinesterase inhibitor.
Atropine is antidotal.

Avoid contamination of food.

5% SEVIN

* Registered trademark of Union Carbide Corp.

GARDEN DUST INSECTICIDE



GENERAL DIRECTIONS FOR USE

Apply 5% Sevin Garden Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 5 to 7-day intervals. For best results, apply with a dust gun. Do not apply in excess.

For use on the following Crops and Ornamentals in Garden: GREENBEANS, LIMABEANS, SNAPBEANS, COWPEAS, BLACK-EYED PEAS, CUCUMBER, MELONS, PUMPKIN, SQUASH, POTATO, TOMATO, EGGPLANT, PEPPER, PEAS, CARROTS, RADISHES, KALE, CABBAGE, BROCCOLI, CAULIFLOWER, GRAPES, STRAWBERRIES, BLUEBERRIES, BLACKBERRIES, BOYSENBERRIES, RASPBERRIES, ROSES, CARNATIONS, GLADIOLUS, CHRYSANTHEMUMS, LILAC, ARBOR VITAE, JUNIPER, HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH, PINES.

Insect Control:
ARMYWORM, BEAN LEAF BEETLE, CUCUMBER BEETLE, CORN EARWORM, COLORADO POTATO BEETLE, CABBAGE CATERPILLAR, EUROPEAN CORN BORER, FLEA BEETLE, HARLEQUIN BUG, JAPANESE BEETLE, LEAFHOPPERS, LIMA BEANS POD BORER, LACE BUGS, MEXICAN BEAN BEETLE, MELON WORM, PICKLEWORM, STINK BUGS, SQUASH BUGS, SPITTLEBUG, TARNISHED PLANT BUG, TOMATO FRUIT WORMS, TOMATO HORNWORM, VELVETBEAN CATERPILLAR, WESTERN BEAN CUTWORMS.

Special Instructions and Pre-Harvest Limitations:

Avoid excess applications on CUCUMBERS, MELONS, PUMPKIN and SQUASH. Some leaf injury may occur if treatments are made when foliage is wet or in the presence of high humidity. For GRAPES, apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers.

There is no limitation on the use of 5% SEVIN GARDEN DUST before harvest for the following crops: GREEN BEANS, LIMA BEANS, SNAP-BEANS, COWPEAS, SWEET CORN, CUCUMBERS, MELONS, PUMPKINS, SQUASH, POTATOES, TOMATOES, EGG-PLANT, PEPPERS, PEAS, GRAPES. Allow three days between dusting and harvesting on the following: CARROTS, RADISHES, KALE, CABBAGE, BROCCOLI, CAULIFLOWER, STRAWBERRIES.

Allow seven days between dusting and harvesting on the following: BLUEBERRIES, BLACKBERRIES, BOYSENBERRIES, RASPBERRIES.

Black Leaf Garden Dust should be applied as often as necessary on small fruit and vegetables listed above for the control of grasshoppers.

When spider mite control is necessary, use a recommended miticide. Do not use Black Leaf Garden Dust containing Sevin while honeybees are visiting the crop.

Do not apply when plants are wet or during a period of high humidity.

FOR USE ON HOUSEHOLD PETS

DOGS, CATS. For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

In and around dog and cat quarters: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.



BLACK LEAF PRODUCTS CO.
Chicago, Ill. 60626

There is no express or implied warranty or representation as to results with this product or effects of its use or as to residues, whether or not in accordance with directions, suggestions, or statements made by the manufacturers of this formulation or of SEVIN or in their behalf, on this label, in advertisements or otherwise. User must assume all risks of use and handling.

BLACK LEAF GARDEN DUST CONTAINING 5% SEVIN

Active Ingredient: Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
Inert Ingredients	95.0%
	100.0%

PLEASE READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

5887-43
12/14/1966

1/1