

ACCEPTED
 MAR 29 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND MITICIDE ACT
 FOR EQUIPMENT REGISTERED
 UNDER 5887-43 SUBJECT
 TO ATTACHED COMMENTS.



* Registered trademark of Union Carbide Corp.

GARDEN DUST

FOR CONTROL OF:



many insects on
 vegetables, small fruits,
 ornamentals.
 Certain pests on
 household pets.

CAUTION:

Keep out of reach of chil-
 dren. (See reverse panel for
 other cautions.)

(The following information will not appear
 on the label:)

- The front panel is 21 sq. inches.
 Caution: Will be printed in 14 point type,
 While "Keep out of reach of children" will
 be printed in 10 point type.
- INERT INGREDIENTS 95.0%
 CCC Diluent supplied by Calcium
 Carbonate Corp., 16.0%
 Pyrox ABB supplied by R.T. Vanderbilt 79.0%
95.0%

SIDE PANEL "A"

SIDE PANEL "B"

CAUTION
 HARMFUL IF INHALED, SWALLOWED,
 OR ABSORBED THROUGH THE SKIN.
 Do Not Breathe Dust. Do Not
 Take Internally. Avoid Prolonged
 or Repeated Breathing of
 Spray.
 Skin Contact May be Harmful.
 Avoid prolonged or repeated con-
 tact with skin. Wash Hands be-
 fore eating.
 Note for Physician: Carbaryl is
 a moderate, reversible, cholin-
 esterase inhibitor. Atropine is
 antidotal.
 Avoid contamination of food.

INSECTICIDE



Distributed by
BLACK LEAF PRODUCTS CO.
 Chicago, Ill. 40626
 U.S.D.A. Reg. No. 5887-43

Buyer assumes all risks
 of use, storage or hand-
 ling of this material, not
 in strict accordance with
 directions given herewith

BLACK LEAF GARDEN DUST CONTAINING 5% SEVIN
 Active Ingredient: Carbaryl (1-naphthyl N-methylcarbamate) 5.0%
 Inert Ingredients 95.0%
 100.0%

PLEASE READ ENTIRE LABEL USE IN STRICT ACCORDANCE
 WITH LABEL DIRECTIONS AND CAUTIONS.

ACCEPTED

3-29-68

UNDER THE FEDERAL INSECTICIDE,
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL
 REGISTERED UNDER NO. 5887-43

BLAR PANEL 5% SEVIN
General Directions for Use

Apply 5% Sevin Garden Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary, generally at 5 to 7 day intervals. For best results, apply with a dust gun. Do not apply in excess.

USE

INSECTS CONTROLLED

SPECIAL INSTRUCTIONS

Crops
 Greenbeans
 Limabeans
 Snapbeans

Mexican bean beetle, Bean leaf beetle, Flea beetle, Japanese beetle, Leafhoppers, Armyworm, Stink Bug, Velvet bean caterpillar, Tarnish plant bug.

Do not apply within 24 hours of harvest.

Cucumber
 Melons
 Pumpkins
 Squash

Pickleworm, Melon worm, Cucumber beetle, Flea beetle, leafhopper, Squash bug,

Avoid excess application, when foliage is tender.

Potato
 Tomato
 Peas
 Eggplant

Colorado potato beetle, Flea beetle, Leafhopper, Evergreen corn borer, Fall armyworm, Lace bug, Tomato fruit worm, Tomato Hornworm, Stink bug, Tarnished plant bug.

ACCEPTED
 3-29-68
UNDER THE FEDERAL INSECTICIDE ACT
 AND REGULATIONS THEREUNDER
 FOR ECONOMIC PEST CONTROL
 REGISTERED UNDER NO. 5887-43

Cabbage
 Broccoli
 Cauliflower
 Lettuce

Flea beetle, Harlequin bug, Armyworm, Cabbage caterpillar, Corn earworm, Small cabbage loopers, Spittlebug, Tarnished plant bug.

Do not apply within 3 days of harvest on cabbage, broccoli and cauliflower, and within 14 days of harvest on lettuce.

Ornamentals

Fls
 Carnations
 Gladioli
 Zinnia

Blister beetle, Flea beetle, Gypsy moth, Japanese beetle, June beetle, Lace bug, leafhopper, leaf roller, Mealy bug, Plant bug, Rose aphid, Tent Caterpillar.

Do not use on Boston Ivy. Do not use when honeybees are visiting plants.

Lilacs
 Magnolia
 Roses
 Evergreens

Bagworm, Boxelder bug, Boxwood leaf miner, Oak leaf miner, Scale insects, Thorn bug, Thrips, Rose Aphid, Rose slug, Japanese beetle, Psyllid.

Small Fruit

Blueberries
 Raspberries
 Blackberries
 Strawberries
 Grapes

Blueberry maggot, Cherry fruitworm, Japanese beetle, Leaf roller, Leafhopper, Raspberry aphid, Meadow spittlebug, Strawberry weevil, Grape berry moth, Japanese beetle, June beetle, Grape leaf skeletonizer, Red banded leaf roller.

Do not apply within 1 day of harvest. Do not use when honeybees are visiting crops.

Citrus
 Grapefruit
 Lemon
 Orange
 Lime

California orange dog, Citrus cutworm, Orange tortrix, Fruit tree leaf roller, Western tussock moth.

Do not apply within 5 days of harvest. Make only 1 application during fruiting season.

House Pets

Dogs,
 Cats

Ticks, Fleas on animal, and in and around animal quarters

Dust animal liberally, avoiding eyes. Comb out 2 hours after treatment. Repeat weekly if needed.