

7-15-71

NET CONTENTS 1



5% SEVIN GARDEN DUST

PLEASE READ ENTIRE LABEL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

GENERAL DIRECTIONS FOR USE

Apply 5% Sevin Garden Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary, generally at 5 to 7 day intervals. For best results, apply with a dust gun. Do not apply in excess

USE	INSECTS CONTROLLED	SPECIAL INSTRUCTIONS
Crops Greenbeans Limabeans Snapbeans	Mexican bean beetle, Bean leaf beetle, Flea beetle, Japanese beetle, Leafhopper, Armyworm, Stink bug, Velvet bean caterpillar, Tarnish ant bug.	Do not apply within 24 hours of harvest.
Cucumbers Melons Pumpkins Squash	Pickworm, Melon worm, Cucumber beetle, Flea beetle, Leafhopper, Squash bug.	Avoid excess application, when foliage is tender.
Potatoes Tomatoes Peas Eggplant	Colorado potato beetle, Flea beetle, Leafhopper, European corn borer, Fall armyworm, Lace bug, Tomato fruit worm, Tomato hornworm, Stink bug, Tarnished plant bug.	
Cabbage Broccoli Cauliflower Lettuce	Flea beetle, Harlequin bug, Armyworm, Cabbage caterpillar, Corn earworm, Small cabbage looper, Spittlebug, Tarnished plant bug.	Do not apply within 3 days of harvest on cabbage, broccoli and cauliflower; and, within 14 days of harvest on lettuce.
Ornamentals Mums Carnations Gladiol Zinnia	Blister beetle, Flea beetle, Gypsy moth, Japanese beetle, June beetle, Lace bug, Leafhopper, Leaf roller, Mealy bug, Plant bug, Rose aphid, Tent caterpillar.	Do not use on Boston Ivy. Do not use when honeybees are visiting plants.
Uliacs Azalea Roses Evergreens	Bagworm, Boxelder bug, Boxwood leaf miner, Oak leaf miner, Scale insects, Thorn bug, Thrip, Rose aphid, Rose slug, Japanese beetle, psyllid.	
Small Fruit Blueberries Raspberries Blackberries Strawberries Grapes	Blueberry maggot, Cherry fruitworm, Japanese beetle, Leaf roller, Leafhopper, Raspberry aphid, Meadow spittlebug, Strawberry weevil, Grape berry moth, Japanese beetle, June beetle, Grape leaf skeletonizer, Red banded leaf roller.	Do not use on Blackberries and Raspberries within 7 days of harvest. Do not apply within 1 day of harvest. Do not use when honeybees are visiting crops.
Citrus Grapefruit Lemon Orange	California orange dog, Citrus cutworm, Orange tortrix, Fruit tree leaf roller, Western tussock moth.	Do not apply within 5 days of harvest. Make only 1 application during

Black Leaf

5% Sevin*

Garden Dust Insecticide

FOR CONTROL OF:
many insects on vegetables, small fruits, ornamentals. Certain pests on household pets.

BLACK LEAF SEVIN* may be used to control insect pests on a wide range of crops and ornamentals. It is an insecticide which will control many insects that are resistant to other pesticides. It may also be used on feed crops.

CAUTION:

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily.

Food utensils such as teaspoons or tablespoons should not be used for food purposes after use with pesticides.

Rinse empty containers thoroughly with water and discard.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to bees.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This container sold by weight and

Black Leaf

5% Sevin*

Garden Dust Insecticide

FOR CONTROL OF:

many insects on vegetables, small fruits, ornamentals. Certain pests on household pets.



BLACK LEAF SEVIN* may be used to control insect pests on a wide range of crops and ornamentals. It is an insecticide which will control many insects that are resistant to other pesticides. It may also be used on feed crops.

CAUTION:

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily.

Food utensils such as teaspoons or tablespoons should not be used for food purposes after use with pesticides.

Rinse empty containers thoroughly with water and discard.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to bees.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.



NET CONTENTS 1 LB.

5% SEVIN GARDEN DUST

PLEASE READ ENTIRE LABEL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

GENERAL DIRECTIONS FOR USE

Apply 5% Sevin Garden Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary, generally at 5 to 7 day intervals. For best results, apply with a dust gun. Do not apply in excess

USE	INSECTS CONTROLLED	SPECIAL INSTRUCTIONS
Crops Greenbeans Limabeans Snapbeans	Mexican bean beetle, Bean leaf beetle, Flea beetle, Japanese beetle, Leafhopper, Armyworm, Stink bug, Velvet bean caterpillar, Tarnish plant bug.	Do not apply within 24 hours of harvest.
Cucumbers Melons Pumpkins Squash	Pickworm, Melon worm, Cucumber beetle, Flea beetle, Leafhopper, Squash bug.	Avoid excess application when foliage is tender.
Potatoes Tomatoes Peas Eggplant	Colorado potato beetle, Flea beetle, Leafhopper, European corn borer, Fall armyworm, Lace bug, Tomato fruit worm, Tomato hornworm, Stink bug, Tarnished plant bug.	
Cabbage Broccoli Cauliflower Lettuce	Flea beetle, Harlequin bug, Armyworm, Cabbage caterpillar, Corn earworm, Small cabbage looper, Spittlebug, Tarnished plant bug.	Do not apply within 3 days of harvest on cabbage, broccoli and cauliflower; and, within 14 days of harvest on lettuce.
Ornamentals Mums Carnations Gladii Zinnia	Blister beetle, Flea beetle, Gypsy moth, Japanese beetle, June beetle, Lace bug, Leafhopper, Leaf roller, Mealy bug, Plant bug, Rose aphid, Tent caterpillar.	Do not use on Boston Ivy. Do not use when honeybees are visiting plants.
Lilacs Azalea Roses Evergreens	Bagworm, Boxelder bug, Boxwood leaf miner, Oak leaf miner, Scale insects, Thorn bug, Thrips, Rose aphid, Rose slug, Japanese beetle, psyllid.	
Small Fruit Blueberries Raspberries Blackberries Strawberries Grapes	Blueberry maggot, Cherry fruitworm, Japanese beetle, Leaf roller, Leafhopper, Raspberry aphid, Meadow spittlebug, Strawberry weevil, Grape berry moth, Japanese beetle, June beetle, Grape leaf skeletonizer, Red banded leaf roller.	Do not use on Blackberries and Raspberries within 7 days of harvest. Do not apply within 1 day of harvest. Do not use when honeybees are visiting crops.
Citrus Grapefruit Lemon Orange Lime	California orange dog, Citrus cutworm, Orange tortrix, Fruit tree leaf roller, Western tussock moth.	Do not apply within 5 days of harvest. Make only 1 application during fruiting season.
House Pets Dogs Cats	Ticks, Fleas on animals, and, in and around animal quarters.	Do not use on puppies and kittens less than 4 weeks old. Dust animal liberally, avoiding eyes. Comb out 2 hours after treatment. Repeat weekly if needed.

DISTRIBUTED BY

BLACK LEAF PRODUCTS CO.
ELGIN, ILLINOIS 60120

ACCEPTED WITH COMMENTS

5% SEVIN GARDEN DUST

PLEASE READ ENTIRE LABEL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

GENERAL DIRECTIONS FOR USE

Apply 5% Sevin Garden Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary, generally at 5 to 7 day intervals. For best results, apply with a dust gun. Do not apply in excess.

USE	INSECTS CONTROLLED	SPECIAL INSTRUCTIONS
<u>Crops</u> Cornbeans Lipbeans Soybeans	Mexican bean beetle, Bean leaf beetle, Flea beetle, Leafhopper, Leaf miner, Armyworm, Stem borer, Yucca bean caterpillar, Tarnish plant bug.	Do not apply within 24 hours of harvest.
<u>Cucumbers</u> Cucumbers Pumpkins Squash	Pickleworm, Melon worm, Cucumber beetle, Flea beetle, Leafhopper, Squash bug.	Avoid excess application, when foliage is tender.
<u>Vegetables</u> Tomatoes Peas Eggplant	Colorado potato beetle, Flea beetle, Leaf miner, Colorado corn worm, Fall armyworm, Lace bug, Tomato fruit worm, Tomato worm, Stem borer, Tarnish plant bug.	
<u>Cabbages</u> Brussels Leekflower Lettuce	Flea beetle, Harlequin bug, Armyworm, Cabbage caterpillar, Corn earworm, Small cabbage looper, Spittlebug, Tarnish plant bug.	Do not apply within 3 days of harvest on cabbage, broccoli and cauliflower; and, within 14 days of harvest on lettuce.
<u>Ornamentals</u> Mosses Carnations Gladiolus Zinnia	Water beetle, Flea beetle, Gypsy moth, Japanese beetle, June beetle, Leaf hopper, Leaf miner, Leaf roller, Milk bug, Plant bug, Rose sawfly, Tent caterpillar.	Do not use on Boston Ivy. Do not use when honeybees are visiting plants.
<u>Lilacs</u> Apples Roses Evergreens	Barkworm, Barklender bug, Boxwood leaf miner, Oak leaf miner, Scale insects, Thorn apple, Tortoise shell beetle, Rose slug, Japanese beetle, psyllid.	
<u>Small Fruit</u> Raspberries Blackberries Strawberries Grapes	Blueberry maggot, Cherry fruitworm, Jack-o-lantern, Leaf miner, Leafhopper, Flea beetle, Strawberry root borer, Strawberry root weevil, Grape berry moth, Japanese beetle, June beetle, Grape leaf skeletonizer, Red-banded leaf roller.	Do not use on Blackberries and Raspberries within 7 days of harvest. Do not apply within 1 day of harvest. Do not use when honeybees are visiting crops.
<u>Citrus</u> Grapefruit Limes Oranges Limes	California orange dog, Citrus cutworm, Orange citrus thrips, Citrus leaf miner, Wax scale, Citrus black fly.	Do not apply within 5 days of harvest. Make only 1 application during fruiting season.
<u>House Pets</u> Dogs Cats	Ticks, Fleas on animals, and, in and around animal quarters.	Do not use on puppies and kittens less than 4 weeks old. Dust animal liberally, avoiding eyes. Comb out 2 hours after treatment.

ANIMAL QUARTERS

It may also be used on

CAUTION:

IF SWALLOWED OR IN-
not take internally. Avoid
dust or spray mist. Avoid
or repeated contact with
h thoroughly after han-
nge contaminated cloth-

ils such as teaspoons or
s should not be used for
ses after use with pesti-

ty containers thoroughly
and discard.

use empty container. De-
y burying with waste or
lay away from smoke or

ct is toxic to bees.

amination of foodstuffs.
amination of feed, feed-
jhs and watering re-

Physician: Carbaryl is a
reversible, cholinesterase
tropic is antidotal.

ner sold by weight and
me. If it does not appear
pened, it is because con-
settled during shipping
ng.