



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

Richard W. Fresh
Consep, Inc., A Subsidiary of Verdant Brands
213 S.W. Columbia St.
Bend, Oregon 97702-1013

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Black Leaf Sevin Garden Dust Insecticide
EPA Registration No. 5887-43
Resubmission Amendment dated February 3, 2000 in Response
to EPA Letter dated 30 November 1999.

APR 18 2000

Dear Mr. Fresh:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

1. In the PHI statement for the crops eggplants, peas, potatoes and tomatoes add the word "not" between the words "Do apply" so that the statement reads "Do not apply within 3 days of harvest on eggplants and tomatoes and 7 days on potatoes".

Submit two copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,
George Tompkins
George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

BLACK LEAF SEVIN® brand carbaryl insecticide GARDEN DUST INSECTICIDE

For Control of Listed Insects on Vegetables, Small Fruits, Ornamentals, Listed Pests on Household Pets

Active Ingredient:	
Carbaryl: (1-Naphthyl N-methylcarbamate).....	5.0%
Other Ingredients:.....	95.0%
Total:	100.0%

Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticide.

EPA Reg. No. 5887-43

EPA Est. No.

Keep Out Of Reach Of Children CAUTION

See Side Panel For Additional Precautionary Statements

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly after handling. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

Wear household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 18 2000

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**
5887-43

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. For best results apply with a dust gun.

CROP USE	INSECT CONTROLLED	SPECIAL INSTRUCTIONS
Green Beans, Lima Beans, Snap Beans	Armyworms, Bean Leaf Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Mexican Bean Beetle, Stink Bugs, Tarnished Plant Bugs, Velvet Bean Caterpillars	Do not apply within 3 days of Harvest for succulents or 21 days For dried.
Cucumbers, Melons, Pumpkins, Squash	Cucumber Beetle, Flea Beetle, Leafhoppers, Melon Worms, Pickleworms, Squash Bug	Do not apply within 3 days of Harvest. Avoid excess application foliage is tender.
Eggplants, Peas, Potatoes, Tomatoes	Colorado Potato Beetle, European Corn Borer, Fall Armyworms, Flea Beetle, Lace Bugs, Leafhoppers, Stink Bug, Tarnished Plant Bug, Tomato Fruitworm, Tomato Hornworm.	Do not apply within 3 days of Harvest for succulent peas and 21 Days for dried peas. Do apply Within 3 days of harvest on Eggplants and tomatoes and 7 days on potatoes.
Broccoli, Cabbage, Lettuce, Cauliflower	Armyworm, Cabbage Caterpillar, Corn Earworm, Flea Beetle, Harliquin Bugs, Small Cabbage Loopers, Spittlebugs, Tarnished Plant Bug.	Do not apply within 3 days of Harvest on cabbage, broccoli, or Cauliflower or within 14 days of Harvest on lettuce
Ornamentals: Carnations, Mums, Gladiolus, Zinnias	Blister Beetle, Flea Beetle, Gypsy Moth, Japanese Beetle, Jun Beetles, Lace Bugs, Leafhoopers, Leaf Rollers, Mealybugs, Plant Bugs, Rose Aphids, Tent Caterpillars	Do not use on Boston Ivy. Do not Use when honeybees are visiting plants.
Azaleas, Evergreens, Lilacs, Roses	Bagworms, Boxelder Bugs, Boxwood Leafminer, Japanese Beetle, Oak Leafminer, Psyllids, Rose Aphids, Rose Slugs, Scale Insects, Thorn Bugs, Thrips	
Small Fruits: Blackberries, Blueberries, Grapes, Raspberries, Strawberries	Blueberry Maggot, Cherry Fruitworms, Grape Berry Moth, Grape Leaf Skeletonizer, Japanese Beetle, June Beetle, Leaf Rollers, Leafhoppers, Meadow Spittlebugs, Raspberry Aphids, Red-banded Leaf Rollers, Strawberry Weevils	Do not apply within 7 days of harvest. Do not apply when honeybees are visiting plants.

PET INSECT CONTROL (Ticks and Fleas)

DOGS: Use this product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Comb out dead ticks a few hours after treatment.

This product should not be applied by children. Do not use on kittens or puppies under 12 weeks of age. Repeat at weekly intervals, if needed.

DOG AND CAT QUARTERS: Dust bedding and dog house using a shaker can or other dust applicator. Force dust into cracks and crevices. Repeat as necessary.

STORAGE AND DISPOSAL

General: Do not contaminate water, food, feed, fertilizer, or other pesticides by use, storage, or disposal. If damaged, leaking, or spilled, sweep up and wrap waste, then dispose of as below. **Storage:** Keep upright, tightly closed in original container in a cool, dry, locked area away from flames, sparks, sunlight, other chemicals, and other sources of ignition, heat, degradation, or contamination. **Disposal:** Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash.

This container sold by weight and not volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with label directions.

For information on this product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide Tele-communications Network at 1-800/858-7378.

If you are not completely satisfied with this product, or for consumer information, call (612) 703-3300 weekdays 9-5 Central Time to arrange a refund for the purchase price or replacement of the product. Proof of purchase is required.

Manufactured by:

Verdant Brands, Inc.

9555 James Ave., South, Suite 200

Bloomington, MN 55431