


1 OZ. NET WT.

Walko

TRADE MARK REG. U.S. PAT. OFF.

TABLETS



DANGER
KEEP OUT
OF REACH
OF CHILDREN

WALKO, a valuable poultry drinking water sanitizer for all poultry to aid in reducing the spread of water-borne disease organisms. For veterinary use only.

Active Ingredient:

Potassium Permanganate 40%

Inert Ingredients: 60%

SEE REAR PANEL FOR FIRST AID AND OTHER PRECAUTIONS:

100 Tablets 5 Gr. Each - Price 75c

EPA REG. NO. 5837-49-AA

DISTRIBUTED BY

BLACK LEAF PRODUCTS CO.

ELGIN, ILLINOIS 60120

00-001

Walko TABLETS

DIRECTIONS FOR USE

For Baby Chicks, Turkeys, Ducks and Geese, 2 Tablets in each quart of drinking water. For Older Birds, 2 to 4 Tablets in each quart of drinking water.

Make fresh solution daily: Oftener if water loses purple color. Stir thoroughly. Do not use in other than non-corrodible type drinking water vessel. Do not mix with milk or feed.

Do not feed chicks until 48 to 60 hours old. Then feed sparingly 4 or 5 times a day.

Discontinue use when birds are receiving any medication or vaccine in their drinking water.

Disease may also be spread by contaminated feed, infected premises, individual contact. Strict sanitary measures should be followed to prevent spread of disease through all sources.

PRECAUTIONS

May be Fatal or Harmful if swallowed
KEEP OUT OF REACH OF CHILDREN

DANGER: Keep out of reach of children. Harmful or fatal if swallowed. Causes severe burns of eye, skin and mucous membranes. Do not get in eyes, on skin, mucous membranes, or on clothing. Avoid contamination of food. Not for drug use. Do not reuse empty container. Discard in a safe place.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Mucosal damage may contraindicate the use of gastric lavage. Measures against respiratory depression and convulsion may be needed.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in accordance with directions given herewith.

\$3.00
\$5.00

LARGE SIZE PACKAGES

1000 TABLETS
2000 TABLETS

ACCEPTED

WALKO TABLETS

080173