

repeat every 7 to 14 days for helminthosporium and rust and every 7 days for

Do not allow to become wet or overheated in storage. This may bring on chemical changes which will affect the film quality.

ACTIVE INGREDIENTS	
A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate	
50% The ingredients are:	
Manganese	16%
Zinc	2%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ S ₄)	82%
INERT INGREDIENTS	20%
TOTAL	100%

* Fore is a registered trademark
of Rohm & Haas Company

EPA REG. NO. 5887-52 AA
Net Weight 8 oz.

Fore

Black
Leaf

Fore

Lawn fungicide

rhizoctonia control

PYTHIUM BLIGHT. Use FORE at 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications at first sign of disease and repeat at 7-day intervals or oftener if conditions are unfavorable for disease development. Continue until disease threat is past.

ANISAE CONTROL ON LAWNS. Use FORE at 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications when disease begins to appear and repeat every 7 days until threat is past.


CELLAR ROT. Use FORE at 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications when disease begins to appear and repeat every 7 days until threat is past. When conditions are unfavorable for disease development, use FORE at 4 oz. per 500 sq. ft. and repeat applications every 7 days until disease threat is past. When conditions are unfavorable for disease development, use FORE at 4 oz. per 500 sq. ft. and repeat applications every 7 days until disease threat is past.

SPOT DISEASE. Use FORE at 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Make applications during the winter months when possible.

For more information on control recommendations for further details on use, see the "CAUTION" label. A complete experimental station is available from the manufacturer.

When using FORE on lawns, use a spray nozzle that will provide adequate coverage.

Will not damage lawns Effective easy-to-use fungicide



CAUTION:
**KEEP OUT OF THE
REACH OF CHILDREN**
See side panel for other cautions

© Fore is a registered trademark
of Black & Leaf Company
EPA REG. NO. 5587-52 AA
Net Weight 8 oz.

bring on chemical changes which will impair the fungicidal effectiveness of Fore and may also generate flammable vapors. Keep container closed when not in use.

Do not reuse container. Destroy when empty.

NOTICE. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S. Patents 2,504,404, 2,504,417 and Patent Pending

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120

PG 720

Fore

Black
Leaf

Fore

Lawn fungicide

rhizoctonia control.

PYTHIUM BLIGHT. Use FORE at 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications at first sign of disease and repeat at 5 day intervals or oftener if conditions are especially favorable for disease development. Continue until disease threat is past.

ALGAE CONTROL ON LAWNS. Use FORE at 3 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications when algae begin to appear and repeat every 7 days until threat is past.

DOLLAR SPOT (SCLEROTINIA). In a protective spray program, use FORE at 3 to 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days. When conditions are especially favorable for dollar spot development, use FORE at 4 oz. per 500 sq. ft. and repeat applications every 7 days until disease is past. A cadmium fungicide may be added to supplement dollar spot control under severe disease conditions.

FUSARIUM SNOW MOLD. Use FORE at 3 to 4 oz. per 500 sq. ft. in sufficient water to provide adequate coverage. Make applications during the winter at a 2 to 6 week interval.

Follow local turf disease control recommendations for further details on timing of applications. Consult State Agricultural Experiment stations or State Extension Service specialist.

When used as directed this package will give coverage up to 2,000 square feet of lawn area.

UNDER THE UNITED STATES
LAW, THIS PRODUCT IS
FOR THE USE OF THE
FEDERAL GOVERNMENT

Will not damage lawns Effective easy-to-use fungicide

Controls: algae, dollar spot, snow mold, brown patch, rust, melting out, slime mold, helminthosporium, copper spot.



bring on chemical changes which will impair the fungicidal effectiveness of Fore and may also generate flammable vapors. Keep container closed when not in use.

Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S. Patents 2,504,404; 2,974,156 and Patent Pending

**DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120**

00-720

101973