

ATTACHMENT 1

A. ACTIVE INGREDIENTS:

Pyrethrins	0.056%
Pine Oil	0.900%
Inert Ingredients:	99.044%

Total	<u>100.000%</u>
-------	-----------------

B. Black Leaf Products  
A Division of Wilbur-Ellis Company  
Buckner, Kentucky 40010

REG # 5887-1-1

PII-11

**DIRECTIONS FOR USE**

**WARNING:** Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130 F. may cause bursting. Never throw container into fire or incinerator.

**CAUTION:** Harmful if swallowed. If used in a food processing or preparation area, foods should be removed or covered during treatment. Food processing surfaces should be covered during treatment or thoroughly cleaned before re-use. Do not treat while food processing is underway.

Shake well before using. Apply this product only as specified on this label.

**GENERAL:** Do not remain in treated areas and ventilate the area after treatment is completed. Remove cap, hold in upright position about 2 feet from object to be sprayed with opening in nozzle pointing in desired direction. Press button to release spray and spray directly on insects where possible.

**FLIES, MOSQUITOS, GNATS:** Close doors and windows of room and remove pets, birds and fish. Thoroughly spray all parts of the room, particularly towards ceiling so as to fill the room with mist. The room should be kept closed 10 to 15 minutes and the fallen insects swept up and destroyed.

**ANTS AND ROACHES:** For best results, spray directly on insects in addition to spraying thoroughly all parts of the room suspected of harboring these pests. Special attention should be paid to cracks and crevices, under sinks, behind stoves and refrigerators and food storage areas. For effective control of ants, spray doorsills, window frames and ant trails and hills. Repeat as necessary.

**SILVERFISH:** For best results, spray directly on insects. For effective control, frequently spray baseboards, doorframes, bookcases and storage areas.

**GARDEN INSECTS:** For use against Japanese beetles, leafhoppers, aphids, exposed thrips. Holding can 1 1/2 feet from plants, spray with one-second bursts, using a sweeping motion. Direct spray on insects. Spray all sides covering plant foliage lightly. Never wet heavily. New growths and young plants should be handled with particular care. Avoid using in direct sunlight or during hottest daylight hours. Repeat when necessary. Do not spray cyclamen, lantana, or maiden-hair fern. Non-staining when used as directed. May be used on: Roses, asters, azaleas, carnations, evergreens, snapdragons, crassula, petunias, rhododendrons, delphiniums, chrysanthemums, iris, African violets, marigolds, grape ivy, English ivy, camellias, euonymus, dahlias.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DISTRIBUTED BY  
**BLACK LEAF PRODUCTS CO.**  
ELGIN, ILLINOIS 60120  
00-450

Not Reviewed. Registrant claims to be in accordance with  
Registration of Pesticide Product Guidance for Phase 2  
Response, Page 2.12, Active to Inert Change in Status.

EPA EST 5887-1L-1

# House & Garden Spray

## Insect killer

Indoors, outdoors,  
garden, and patio

ACTIVE INGREDIENTS	BY WEIGHT
PYRETHRINS	0.25
N-OCTYL BICYCLOHEPTENE DICARBOXIMIDE	0.50
TECHNICAL PIPERONYL BUTOXIDES	0.50
PETROLEUM DISTILLATE	8.75
INERT INGREDIENTS	90.00
	100.00

**CAUTION:**  
KEEP OUT OF THE  
REACH OF CHILDREN

CONTENTS UNDER  
PRESSURE

SEE REAR PANEL FOR OTHER CAUTIONS

EQUIVALE  
(6-PROPYL  
RELATED

EPA R

No

A

B