



# Lawn


## weed killer



**ACCEPTED**

JAN 12 1973

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION SUBJECT TO THE FOLLOWING CONDITIONS:



### Kills Dandelions weed killer

**ACTIVE INGREDIENTS:**  
 Dimethylamine Salt of Dicamba (3, 6-Dichloro-o-anisic Acid)\*\* 3.18%  
 Dimethylamine Salts of related acids .52%  
 Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid\*\* 12.72%  
**INERT INGREDIENTS** 83.58%  
 TOTAL 100.00%

\*Equivalent to 2.64% 3,6-Dichloro-o-anisic acid (Dicamba) or 0.23 pounds per gallon  
 \*\*Equivalent to 10.57% 2,4-Dichlorophenoxyacetic acid or 0.92 pounds per gallon.

32 FL. OZS. (1 QUART)  
 E.P.A.Reg. No. 5887 68

EFFECTIVE FOR THE CONTROL OF  
**DANDELIONS**  
**CLOVER IN LAWNS**  
**POISON IVY, POISON OAK**  
 CONTAINS 2,4-D FOR DANDELIONS  
 DICAMBA FOR GENERAL WEED CONTROL

**WARNING:**  
 KEEP OUT OF THE REACH OF CHILDREN.  
 See other panel for other cautions

MADE IN U.S.A.



### Lawn WEED KILLER GENERAL DIRECTIONS FOR USE

Lawn Weed Killer contains both 2,4-D and Dicamba for effective control of a broad range of weeds infesting lawns.

**LAWN WEEDS:** To kill dandelion, wild onion, pennywort, plantain, heal-all and similar weeds, use 8 ounces in 6 gallons of water and apply with sprayer or sprinkling can to 2,500 square feet of lawn. (For smaller areas use 2 teaspoonsful in 2 quarts of water applied to 100 square feet.)  
 Do not apply to areas underlaid by roots of desirable trees and shrubs.

To kill weeds such as thistles, bindweed, dock and ragweed, use 8 ounces in 6 gallons of water (or 2 teaspoonsful in 2 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary.

To kill undesirable plants such as poison ivy and poison oak, use 8 teaspoonsful in 2 quarts of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard to kill weeds such as poison ivy, poison oak, English daisy may require a fall application.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith

**WARNING:** Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

**IMPORTANT:** Avoid drift of spray (coarse sprays are less likely to drift) on vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides. Do not use on centipede grass or dichondra nor on lawns or turf where desirable bent grass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time. To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period. When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

**BLACK LEAF LAWN WEED KILLER** is effective for the control of dandelion, chicory, clover, dock, dog fennel, English daisy, Frenchweed, knotweed erect, knotweed prostrate, ground ivy, hawkweed, heal-all, henbit, lambsquarter, mallow, morning glory, chickweed (mouse ear-common-stitchwort); mustard, oxalis, pennywort, peppergrass, pigweed, plantain, poison ivy, poison oak, Canada thistle, burdock, jimson weed, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, wild aster, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, and many other broadleaf weeds.

Distributed by  
**BLACK LEAF PRODUCTS CO., ELGIN, ILLINOIS 60120**  
 00 003

This product is licensed under one or more of the following United States patents: 2,390,941 2,394,916 2,396,513 2,412,510 2,453,983 2,472,347 U.S. Reg. Pat. No. 23,115