

**WARNING: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**WARNING: HARMFUL IF SWALLOWED:** Avoid breathing of mist or contact with skin. In case of contact, wash with soap and water. Keep out of reach of children.

This product is toxic to fish. Keep out of all bodies of water. This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

**DO NOT USE** with materials containing sulphur or other substances incompatible with oil. **DO NOT USE** on crops within 14 days of application of any material containing sulphur.

**DO NOT** contaminate water by cleaning of equipment or disposal of wastes.

**DO NOT** contaminate feed or foodstuffs.

**DO NOT** re-use container for any purpose. Do not use in any manner not described on this label.

**NOTICE**

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

**DISTRIBUTED BY**  
**BLACK LEAF PRODUCTS COMPANY**  
ELGIN, ILLINOIS 60120  
00-212

**Black  
Leaf**

**25%**

**Methoxychlor**

**mosquito-insect spray**

**KILLS Mosquitoes,  
Japanese Beetles and  
other listed insects  
on fruit trees, ornamentals,  
flowers and vegetables.  
KILLS Cankerworms (Inch Worms)  
on shade trees.**

**DIRECTIONS FOR USE OF BLACK LEAF 25% METHOXYCHLOR**

**LEAF FEEDING INSECTS ON FLOWERS, SHADE TREES AND ORNAMENTALS:** Mix 4-6 fluid ounces of concentrate per 6 gallons of water. (1 1/3 to 2 tablespoonfuls per gallon). Apply to thoroughly cover all plant surfaces when small larvae begin feeding.

**ON SHADE TREES ORNAMENTALS:** Controls Cankerworms (Inch Worms), Gypsy Moth, Juniper Webworm.

**ON FLOWERS:** Apply to Roses, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemum, Coleus, Dahlias, Delphiniums, Geraniums, Gladioli, Marigolds, Nasturtiums, Snap Dragons, and Zinnias to control Japanese Beetles, Leaf Hoppers, Lacebugs, Blister Beetles, Cucumber Beetles, Flea Beetles, Rose Chafers, and Rose Slugs.

Consult your Agriculture Extension Service for best time to spray in your area.

**DO NOT** spray Chinese Elms, Japanese Maples, Red Maples and Redbud.

**DO NOT** spray when temperature is above 85°F.

**FRUITS:** Apply only to Apples, Pears, Peaches, Apricots, Cherries and Plums. Mix 4 Fluid Ounces of concentrate per 6 gallons of water, spray all surfaces of foliage and fruit. Begin at petal fall and repeat at 7 to 14 day intervals as needed to control Apple Maggot, Cherry Fruit Worm, Codling Moth, Leaf Hoppers, Plum Curculio, Tent Caterpillar and Japanese Beetle. Do not apply later than 7 days before harvest on Apples, Cherries, Pears, Plums or later than 21 days of harvest on Peaches and Apricots.

Apply the same concentration to Strawberries, Blueberries, Raspberries, Blackberries, Currants, Gooseberries, and Grapes. To control Grape Berry Moth, Grape Leaf Skeletonizer, Rose Chafer, Leaf Hopper, Spittlebug, and Strawberry Weevil. **DO NOT** apply within 14 days of picking. Wash residue off all crops before using.

**VEGETABLES:** Mix 4-6 Fluid Ounces of concentrate per 6 gallons of water. Apply to Tomatoes, Beans, Cabbages, Cauliflower, eggplants, peas, peppers, and potatoes to control Bean Leaf Beetles, Blister Beetles, Colorado Potato Beetles, Japanese Beetles, Leaf Hoppers, Mexican Bean Beetles and Lace Bugs. **DO NOT** apply within 7 days of picking. Wash residue off all crops before using.

**AREA MOSQUITO SPRAY:** To aid in the control of Mosquitos, Gnats, and certain other flying insects around home grounds, patios, and picnic areas. Mix 2 Fluid Ounces of concentrate in 3 gallons of water. Apply as a coarse spray to resting surfaces such as grass, shrubs and tree trunks and branches to a height of 8 to 10 feet. Thoroughly wet plants but do not apply to point of run off on the soil. Do not spray patio surfaces. Keep children and pets off treated areas until spray dries. May also be applied to the same areas with a fog or mist applicator using 1 part concentrate to 4 parts of water. Best results are obtained by applying in early morning or night. Do not apply more than 13 Fluid Ounces of concentrate (0.2 pounds of actual Methoxychlor per acre). Repeat as necessary.

5-63-22  
5-63-74

DO NOT USE, POUR, SPILL  
NEAR HEAT OR OPEN FLAME.

HARMFUL IF SWALLOWED:  
thing of mist or contact with  
se of contact, wash with soap  
Keep out of reach of children.

ct is toxic to fish. Keep out of  
of water. This product is toxic  
d should not be applied when  
ctively visiting the area. Do not  
low to drift to areas occupied  
ected humans or beneficial

USE with materials containing  
other substances incompatible  
DO NOT USE on crops within 14  
application of any material con-  
phur.

ontaminate water by cleaning of  
or disposal of wastes.

ontaminate feed or foodstuffs.

-use container for any purpose.  
e in any manner not described  
el.

#### NOTICE

imes all risks of use, storage,  
g of this material not in strict  
with directions given herewith.

DISTRIBUTED BY  
EAF PRODUCTS COMPANY  
GIN, ILLINOIS 60120

00-212

**Black  
Leaf**

**25%**

**Methoxychlor**

**mosquito-insect spray**

**ACTIVE INGREDIENTS** By Weight  
\*Methoxychlor, Technical 25.0%  
Xylene 72.0%  
**INERT INGREDIENTS** 3.0%  
**Total 100.0%**  
\*25% Methoxychlor, technical is equi-  
valent to 22% 2,2-bis (p-Methoxyphenyl)-  
1,1,1-Trichloroethane and 3% other iso-  
mers and reaction products.

## WARNING

**KEEP OUT OF THE  
REACH OF CHILDREN**

See side panel for other warnings.

**EPA REG.NO.5887-79-AA**

**NET CONTENTS 16 FL. OZ. (1 PINT)**

**DIRECTIONS FOR USE OF  
LEAF FEEDING INSECTS ON FLOWERS:**  
4-6 fluid ounces of concentrate per  
gallon). Apply to thoroughly  
feeding.

**ON SHADE TREES ORNAMENTAL**  
Moth, Juniper Webworm.

**ON FLOWERS:** Apply to Roses, As-  
Chrysanthemum, Coleus, Dahlias,  
Nasturtiums, Snap Dragons, and  
pers, Lacebugs, Blister Beetles, Cu,  
Rose Slugs.

Consult your Agriculture Extension  
DO NOT spray Chinese Elms, Jap  
DO NOT spray when temperature

**FRUITS:** Apply only to Apples, Pe  
4 Fluid Ounces of concentrate per  
and fruit. Begin at petal fall and r  
trol Apple Maggot, Cherry Fruit  
culio, Tent Caterpillar and Japane  
harvest on Apples, Cherries, Pear  
Peaches and Apricots.

Apply the same concentration to  
berries, Currants, Gooseberries, a  
Leaf Skeletonizer, Rose Chaf  
DO NOT apply within 14 days of

**VEGETABLES:** Mix 4-6 Fluid Ounc  
to tomatoes, beans, cabbages, ca  
to control Bean Leaf Beetles, Bli  
Beetles, Leaf Hoppers, Mexican B  
7 days of picking. Wash residue of  
**AREA MOSQUITO SPRAY:** To aid  
other flying insects around home  
Ounces of concentrate in 3 gallo  
surfaces such as grass, shrubs at  
10 feet. Thoroughly wet plants b  
Do not spray patio surfaces. Kee  
dry. May also be applied to the  
1 part concentrate to 4 parts of  
can be used at night. Do not a  
(0.1) pounds of 2,2-bis (p-Methoxychl