



Reg. T/M of Union Carbide Corp.

**Controls Listed Insects—
including Chinch Bugs**



**Can Be Used
On: Lawns
Ornamentals
Vegetables
Fruits
Citrus Fruit**

CAUTION:
**KEEP OUT OF THE
REACH OF CHILDREN**
See side panel for other cautions

BLACK LEAF SEVIN* may be used to control insect pests on a wide range of crops and ornamentals. It is an insecticide which will control many insects that are resistant to other pesticides.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily.

Food utensils such as teaspoons or tablespoons should not be used for food purposes after use with pesticides.

Rinse empty containers thoroughly with water and discard.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to bees.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This container sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping and handling.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

Please follow directions and cautions carefully before each use. The chart is based on tablespoonsful added to each gallon of water.

USE	INSECTS CONTROLLED	DIRECTIONS
Ornamentals Carnations Gladioli Zinnia Mums Lilacs Azaleas Roses Evergreens	Apple aphid, Bagworm, Birch leaf miner, Blister beetle, Boxelder bug, Elm leaf aphid, Flea beetle, Japanese beetle, June beetle, Lace bug, Leafhopper, Leaf roller, Mealy bug, Orange tortrix, Periodical cicada (locust), Plant bug, Rose aphid and slug, Scale insects, Tent caterpillars.	Mix 2 tablespoons per gallon of water and spray entire plant thoroughly. Repeat applications weekly. Do not use on Boston Ivy.
Vegetables Beans Cabbage Broccoli Lettuce Carrots Cucumbers Squash Tomatoes	Mexican bean beetle, Flea beetle, Japanese beetle, Leafhopper, Velvet bean caterpillar, Armyworm, Corn earworm, Stink bug, Tarnished plant bug, Hadequin bug, Cabbage caterpillar, Six-spotted leafhopper, Corn rootworm adults, Sap beetle, Bean leaf beetle, Pickleworm, Lace bugs, Tomato Fruitworms and hornworms.	Spray sufficient gallonage to obtain full coverage of plants using 2 to 4 tablespoons per gallon of water. Keep spray well agitated. Repeat weekly and as damage appears. Do not apply within 3 days of harvest on Broccoli and Cabbage, and within 14 days of harvest on Lettuce.
Fruits Apples Pears Peaches Apricots Strawberries Cherries Plums	Apple mealybug, Aphids, Leafhoppers, Apple maggot, Bagworm, Eastern tent caterpillar, Bud moth, and fruit tree leaf rollers, Fruitworms, Japanese beetle, Pear psylla, Plum curculio, Rosy and woolly apple aphids, Rust mite, Blister mite, Tarnished plant bug, Tentiform leaf miners, Scale, Oriental fruit moth, European earwig, June beetles, Pear and Peach twig borer, Tussock moth, Meadow spittlebug, Strawberry weevil, Black Cherry aphid, Mealy plum aphid, Cherry fruit fly, Leafhopper, Cutworm, Fireworm, Filbert aphid, Filbert moth, and leaf roller.	Mix 2 tablespoons of powder per gallon of water. Keep mixture well agitated. Spray thoroughly on entire tree or plant. Repeat weekly or as soon as pests reappear. Do not apply within 3 days of harvest on apricots, within one day of harvest on apples, pears, strawberries, cherries and plums.
Citrus Fruit Oranges Lemons Grapefruit Lime	Citrus cutworm, California orange dog, Fruit tree leaf roller, Orange tortrix, Western tussock moth, Scale pests.	Mix 2 tablespoons per gallon of water. Spray trees and branches thoroughly. Repeat soon as pests reappear. Do not apply within 5 days of harvest on all citrus.
Lawns Ornamental Turf Patio	Mosquitoes, Ants, Chinch bugs, Fall armyworms, Leafhoppers, Sod webworms, Fleas.	15 gallons of spray to cover 500 sq. ft. using 1 tablespoon of powder per gal. Spray sufficiently to insure penetration of turf. Repeat applications every 2 weeks.

**DISTRIBUTED BY
BLACK LEAF PRODUCTS CO.**

ELGIN, ILLINOIS 60120

00 610

