

Lawn Weed Killer

AC
Ja
UNDER
FUNGIC
FOR EC
ED UND

KILLS dandelions and weeds

... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..



WARNING: KEEP OUT OF THE REACH OF CHILDREN

NET CONTENTS 32 FL. OZ. (1 QUART) EPA REG. NO. 5887-92-AA



Lawn Weed Killer

ACCEPTED
January 16, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTGENIC ACT
FOR PERSISTENT POISON REGISTERED
UNDER NO. 5887-92

Kills dandelions and weeds

ALSO KILLS CLOVER POISON IVY POISON
OAK AND WOOLLY UNDESIRABLE WEEDS
CONTAINS 2,4-D AND SILVEX

ACTIVE INGREDIENTS:
*Dimethylamine salt of 2,4-D
(2,4,5-Trichlorophenoxyacetic acid)
4.00%
**Dimethylamine salt of 2,4-D
Dichlorophenoxyacetic acid
10.97%
INERT INGREDIENTS 87.03%
TOTAL 100.00%

Equivalent to 3.43% Silvex
(2,4,5-Trichlorophenoxypro-
panoic acid) or 30 pounds per
gallon
or 10.97% 2,4-D
or 10.97% 2,4-D and 0.92
pounds per gallon



WARNING: KEEP OUT OF THE REACH OF CHILDREN
See rear panel for additional warning

NET CONTENTS 32 FL. OZ. (1 QUART) EPA REG. NO. 5887-92-AA

HOW TO USE

Apply 2.0 to 4.0 quarts of solution per 1,000 sq. ft. of lawn. For best results, apply to lawn when weeds are in full leaf. For best results, apply to lawn when weeds are in full leaf. For best results, apply to lawn when weeds are in full leaf.

For best results, apply to lawn when weeds are in full leaf. For best results, apply to lawn when weeds are in full leaf. For best results, apply to lawn when weeds are in full leaf.

WARNING: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water. For eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides. Do not use on centipede grass, St. Augustine or cordgrass nor on lawns or turf where desirable bent grass or clover are present. Newly seeded lawns may be treated after they have been mowed the second time. To reseed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period. When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

BLACK LEAF LAWN WEED KILLER is effective for the control of dandelion, chicory, clover, dock, dog fennel, English daisy, knotweed, knotweed erect, knotweed prostrate, ground ivy, horseweed, leafy spurge, henbit, lamb's quarter, mallow, morning glory, chickweed (mouse ear common-stitchwort), mustard, oxalis, pennywort, peppergrass, pigweed, plantain, poison ivy, poison oak, Canada thistle, burdock, jimson weed, purslane, ragweed, smartweed, sleep sorrel, spotted spurge, wild aster, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, timothy and yarrow.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
00-803

