

**DIRECTIONS FOR USE OF THURICIDE®**

**GENERAL:** Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Good underleaf coverage is essential since the insect must ingest Black Leaf THURICIDE®. After eating Black Leaf THURICIDE®, insects stop feeding but may otherwise appear to be unaffected for several days. An adhesive or deposit builder may help where heavy dew or rains are encountered.

Store in a cool place. Protect from freezing.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy or discard in a safe place.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

THURICIDE® Reg. Trade Mark of International Minerals and Chemicals Corporation.

DISTRIBUTED BY  
**BLACK LEAF PRODUCTS COMPANY**  
ELGIN, ILLINOIS 60120

00-141



# Thuricide®

a microbial insecticide

**FOR USE ON ORNAMENTALS, SHADE TREES, and VEGETABLES.** Effective for the control of Spring and Fall Cankerworms (Inch Worms), Gypsy Moths (NE), and other chewing worms specified in the directions on this label.



**DIRECTIONS FOR USE OF BLACK LEAF THURICIDE**

Shake or stir suspension before use. Pour recommended amount of BLACK LEAF THURICIDE onto surface of water in nearly-filled spray tank. Maintain agitation. Do not allow diluted spray to stand in tank for more than 12 hours. Combinations with chemical insecticides, fungicides, spreader-stickers in the tank are generally not deleterious to BLACK LEAF THURICIDE if used promptly.

| FOR USE ON  | CONTROLS THESE PESTS   | RECOMMENDED USAGE  |
|---|--|--|
| Shade trees   | Cankerworms (Spring and Fall)<br>Fall Webworm<br>Tent Caterpillar<br>Gypsy Moth (NE) | Use 2 fl. oz. of THURICIDE in 3 gallons of water (4 teaspoonsful per gallon). For power sprayers use 1 to 2 quarts in 100 gallons of water per acre. Apply when worms begin to hatch. Cover foliage thoroughly. Repeat application after 14 days if necessary. |
| Fruit   |  |  |
| Grapes  | Grape Leaf Folder  | Mix 1½ fl. oz. of Black Leaf THURICIDE to 3 gallons of water or 1 tablespoon per gallon. Apply to all foliage covering an area up to 1,000 sq. ft. Apply as soon as pests are noticed and repeat weekly as required for control of pests.                      |
| Oranges (Calif. and Ariz.)                                      | Fruit Tree Leafroller<br>Orange dog  |  |
| Vegetables  |  |  |
| Broccoli<br>Cabbage<br>Cauliflower<br>Collards<br>Kale          | Cabbage Looper<br>Imported Cabbage worm  | Use 1½ fl. oz. in 3 gallons or 1 tablespoon per gallon in a hand sprayer and apply to all foliage in an area up to 1,000 sq. ft.   |
| Celery<br>Lettuce<br>Cucumbers<br>Melons<br>Potatoes<br>Spinach | Cabbage Looper   |  |
| Tomatoes  | Cabbage Looper<br>Tomato Hornworm  |  |

When used as directed this package will make up to 16 gallons of spray.

5887-9L

12-13-72

1 of 2

011111

1077

ACCEPTED  
DEC 13 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTE-  
RED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

5887-96

2012