2lb. Sevin®

and as Vegetable Garden Spray, Controls Listed Lawn Garden and Ornamental Plant Insects.

ACTIVE INGREDIENT: Carbaryi (1-naphthyl-N-methylcarbamate) WERT INGREDIENTS		
Total	100.0%	
CAUTION		

KEEP OUT OF THE REACH OF CHILDREN

CAUTION:

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust or spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Avoid contamination of food

This product is highly toxic to pollinating

bees exposed to direct treatment or re-

sidues on crops. Protective Information

may be obtained from your Coorperative Agriculture Extension Service.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe

NOTICE: Buyer assumes all risks of use. storage, or handling of this material not in

strict accordance with directions given

herewith. Protect from temperatures below

32 F. Keep cover on opened container to

Sevin' Trade Mark of the Union Carbide Corp. for the active ingredient. 1-

naphthyl-N-methylcarbamate U.S. Pat. No.

DISTRIBUTED BY **BLACK LEAF PRODUCTS COMPANY**

ELGIN. ILLINOIS 60120

Contains 2 pounds of Sevin per gation

prevent drying.

2.903.478.

and keep out of reach of children.

EPA REG. NO. 5887-102 AA

NET CONTENTS 16 FL. OZ. (1 PINT)

DIRECTIONS FOR USE OF LIQUID SEVIN SHAKE WELL BEFORE USING LIQUID SEVIN is an extremely fine dispersion of Sevin in water. Pour

recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixting and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.

PRECAUTIONS: Avoid spray drifting to adjoining food or forage crops in bloom. For protection of honey bees, avoid unnecessary use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Refer to State or Federal Agricultural Authorities for more information on the control of the pests listed. Sevin is compatible with common insecticides, fungicides and miticides, but not with lime or other highly alkaline materials. LAWN SPRAY: For ants, chinch bugs, earwigs, fleas, fall armyworm, mosquitoes, sod webworms (lawn moths) and millipedes. For best results treat after cutting and when the ground is moist from rain or irrigation using ½ pint in 15 to 20 gallons of water to each 500 square feet of established lawn. Apply with hose end or pressure sprayer with ample water to insure turf penetration. Repeat in 2 or 3 weeks if neces-

sary.
ORNAMENTAL PLANT SPRAY: For flowers, shrubs and shade trees including roses, azaleas, mums, lilac, arborvitae, juniper, pine, birch, dogwood, oak and maple. (Do not use on Boston Ivy). Apply spray mixture containing 4 teaspoonfuls per gallon of water (8 fl. oz. to 12 gallons or I quart to 50 gallons) when insects or damage appear. Repeat at I quart to 50 gallons) when insects or damage appear, Repeat at weekly intervals or as necessary for such insects as bagworm, leaf miners, elm leaf beetle, gypsy moth, Japanese beetle, lacebug, rose slug, tent caterpillars, leaf rollers, scale insects and periodical Cicada. VEGETABLE GARDEN SPRAY: Mix 2 oz. in 3 gallons of water (2 quarts in sufficient water for 1 acre.) Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with directions!

BEANS (green lima and snap)—Japanese and Mexican Bean Beetle, Tarnished Plant Bug. TOMATO, EGGPLANT & PEPPER—Leaf hopper, lacebug, tomato fruit worm. PUMPKIN, SQUASH, CUCUMBER, MELON—Flea beetle, melonworms, cucumber beetle, squash bug. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity, Do not use on watermelons in Florida.

Kills chinch bugs-gypsy moths

USE ON: Lawns, Flowers, Shrubs and Shade Trees

CAUITUN See side ganel for

ACCEPTED

FEB 1 5 1974 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER. EQ UNDER NO. 5817-10 SUBJECT TO ATTACHED COMMENTS.