

DEC 13 1991

Ms. Joanne Pratt
Wilbur-Ellis company
P.O. Box 9518
Fresno, California 93792

Dear Ms. Pratt:

Subject: Revised Labeling/ Confidential Statement of Formula
Black Leaf Liquid Flowable 2 lb. Sevin
EPA Registration No. 5887-102
Your Application Dated September 5, 1991

The revised labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable. A stamped copy is enclosed for your records.


The Confidential Statement of Formula dated August 15, 1991 is acceptable and will be placed on file for your product.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Amended
Label



Black Leaf (R)

Liquid Sevin (R) *
Brand Carbaryl Insecticide

Kills Ants-Gypsy Moths
Let's you control lawn, vegetable or flower garden, and
ornamental plants from marauding insects!

ACTIVE INGREDIENT:

Carbaryl + (1-naphthyl- N-methylcarbamate).....	21.3%
INERT INGREDIENTS.....	78.7%
TOTAL	100.0%

*SEVIN is a registered trademark of Rhone-Poulenc for
carbaryl insecticide.
+Equivalent to 2 lb carbaryl per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional precautionary statements on right panel.

Reg. No. 5887-102
EPA Est. No. 70-KY-1
NET 16 FL OZ (1 PT) 473 mL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid breathing spray
mist or vapors. Avoid contact with eyes, skin, or clothing.
Wash thoroughly after handling. Avoid contamination of food.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible
cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine
invertebrates. Do not apply directly to water or wetlands.
Do not contaminate water by cleaning of equipment or disposal
of wastes.


This product is highly toxic to bees exposed to direct
treatment or residues on blooming crops or weeds. Do not
apply this product, or allow it to drift, to blooming crops
or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a
manner inconsistent with its labeling.

SHAKE WELL BEFORE USING

This product is an extremely fine dispersion of carbaryl in
water. Pour recommended amount of material into nearly full
spray tank, or pre-mix material in another container half-
full of water and pour mixture into spray tank. Add balance



of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces. Avoid spray drifting to adjoining food or forage crops in bloom.

Carbaryl is compatible with common insecticides, fungicides, and miticides, but not with lime or other highly alkaline materials.

LAWN SPRAY: For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Millipedes, Mosquitoes, and Sod Webworms (Lawn Moths). For best results, treat after cutting and when the ground is moist from rain or irrigation, using 1/2 pt in 15 to 20 gal. of water for each 500 sq ft of established lawn. Apply with hose-end or pressure sprayer, with ample water to insure turf penetration. Repeat in 2 or 3 weeks if necessary.

ORNAMENTAL PLANT SPRAY: For flowers, shade trees, and shrubs, including Arborvitae, Azaleas, Birches, Dogwoods, Junipers, Lilacs, Maples, Mums, Oaks, Pines, and Roses. Do not use on Boston Ivies. Apply spray mixture containing 4 teaspoonfuls per gal. of water (8 fl oz to 12 gal) when insects or damage appear. Repeat at weekly intervals, or as necessary, for such insects as Bagworms, Elm Leaf Beetles, Gypsy Moths, Japanese Beetles, Lace Bugs, Leaf Miners, Leaf Rollers, Periodical Cicadas, Rose Slugs, Scale Insects, and Tent Caterpillars.

VEGETABLE GARDEN SPRAY: Mix 2 fl oz in 3 gal of water. Spray when insects or damage appear, and repeat at 7 to 10 day intervals or when necessary. No post-treatment time limitations for the following host plants when material is applied in accordance with directions:

Beans (Green, Lima, and Snap)- Japanese Bean Beetles, Mexican Bean Beetles, and Tarnished Plant Bugs.


Eggplants, Peppers, and Tomatoes- Lace Bugs, Leafhoppers, and Tomato Fruit Worms.

Cucumbers, Melons, Pumpkins, and Squashes - Cucumber Beetles, Flea Beetles, Melonworms, and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet, or in the presence of prolonged high humidity. Do not use on Watermelons in Florida.

STORAGE AND DISPOSAL

GENERAL: Do not contaminate water, food, feed, or other pesticide containers, storage, or disposal. If damaged, leaking, or spilled, soak up and wrap waste, then dispose of as below.

STORAGE: Keep in upright, tightly closed original container in a cool, dry, locked area away from flames, sparks, sunlight, other chemicals, or other sources of ignition,



BlackLeaf Liquid Sevin (Bleaf\Liqsevin)

Page 3

heat, degradation, or contamination. Keep above 32F (0C); if allowed to freeze, thaw and shake thoroughly before using.

DISPOSAL: Do not reuse empty container. Rinse original container thoroughly, then securely wrap in several layers of newspaper and discard in trash.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

UPC CODE# 76342 00632

DISTRIBUTED BY

BLACK LEAF PRODUCTS COMPANY, A Division of Wilbur-Ellis
Company

BUCKNER, KENTUCKY 40010

B-891 Made in U.S.A.

