	-			····· ·····					
, ,	ЪW	04	5887-1	02	6/18/98	2	page	175	
	BL	ACK LEAF		ABLE 2 LB. S	SEVIN [®] brand ca	arbaryl inse	cticide		
		Kills Ants - Gypsy Moths							
	Lets y	Lets you control lawn, vegetable or flower garden, and ornamental plants from marauding insects!							
	Carba	gredients:	hthyl N-methylca	•					
	Equivalent to 2 lb. carbaryl per gallon Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticities CEPTED								
	EPA Re	eg. No. 588	37-102	EP	A Est. No. 769-	GA-1	JUN 18	998	
	KEEP OUT OF REACH OF CHILDREN CAUTION					N Fun as s regi	Under the Federal Insecticide, Fundicide, and Rodenticide Act, as amended, for the peakide registered under EPA Reg. No. 5887-102		
	See Side Panel For Additional Precautionary Stateme							r • • • • •	
		i	PRECAUTIO Hazards To Hum	ONARY STA nans and Do CAUTION		S) () () () () () () () () () (
	contact contam Note To	with eyes, ination of f	n: Carbaryl is a m	. Wash thor	oughly after har	ndling. Av	oid		
			ENVIRON	IMENTAL H	AZARDS				
	This po	cticido in o	véromoly toyic to	aquatic and	ostuprine inver	tohratos	Do pot		

٩

ť

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product, or allow it to drift, to blooming crops or weeds if bees are visiting the treatment area.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering. För trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper sand lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

SHAKE WELL BEFORE USING

This product is an extremely fine dispersion of carbaryl in water. Pour recommended amount of material into nearly full spray tank, or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces. Avoid spray drifting to adjoining food or forage crops in bloom.

Carbaryl is compatible with common insecticides, fungicides, and miticides, but not with lime or other highly alkaline materials.

Lawn Spray: For Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Millipedes, Mosquitoes, and Sod Webworms (Lawn Moths). For best results, treat after cutting and when the ground is moist from rain or irrigation, using 1/2 pint in 15 to 20 gallons of water for each 500 sq. ft. of established lawn. Apply with hose-end or pressure sprayer, with ample water to insure turf penetration. Repeat in 2 or 3 weeks if necessary.

Ornamental Plant Spray: For flowers, shade trees and shrubs, including Arborvitae, Azaleas, Birches, Dogwoods, Junipers, Lilacs, Maples, Mums, Oaks, Pines, and Roses. Do not use on Boston Ivies. Apply spray mixture containing

2

295

4 teaspoonfuls per gallon of water (8 fl. oz. to 12 gal.) when insects or damage appear, for such insects as Bagworms, Elm Leaf Beetles, Gypsy Moths, Japanese Beetles, Lace Bugs, Leafminers, Leaf Rollers, Periodical Cicadas, Rose Slugs, Scale Insects, and Tent Caterpillars.

Vegetable Garden Spray: Mix 2 fl. oz. in 3 gallons of water. Spray when insects or damage appear. Do not apply within 3 days of harvest.

Beans (green, lima and snap): Japanese Bean Beetles. Mexican Bean Beetles, and Tarnished Plant Bugs.

Eggplants, Peppers, and Tomatoes: lace Bugs, Leafhoppers, and Tomato Fruit Worms.

Cucumbers, Melons, Pumpkins, and Squashes: Cucumber Beetles, Flea Beetles, Melonworms, and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet, or in the presence of prolonged high humidity. Do not use on Watermelons in Florida.

STORAGE AND DISPOSAL

General: Do not contaminate water, food, feed, or other pesticides by use, storage, or disposal. If damaged, leaking, or spillec, soak up and wrap waste, then dispose of as below.

Storage: Keep upright, tightly closed original container in a cool, dry, locked area away from flames, sparks, sunlight, other chemicals, or other sources of ignition, heat, degradation, or contamination. Keep above 32° F (0° C); if allowed to freeze, thaw and shake thoroughly before using **Disposal:** Do not reuse empty container. Rinse original container thoroughly;

then securely wrap in several layers of newspaper and discard in trash.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Net Weight:

Manufactured for: Black Leaf Products Louisville, KY 40222

5/98 EPA stamped label 4/14/98

ATTACHED SPRAYER DIRECTIONS FOR USE

Lawn Spray: For Ants, Chinch Bugs. Earwigs, Fall Armyworms, Fleas, Millipedes, Mosquitoes, and Sod Webworms (Lawn Moths). For best results, treat after cutting and when the ground is moist from rain or irrigation. Activate attached sprayer using 8 ounces of this product to treat 1,333 sq. ft. of established lawn. Apply with ample water to insure turf penetration. Repeat in 2 or 3 weeks if necessary.

Ornamental Plant Spray: Use for insect control on flowers, shade trees and shrubs, including Arborvitae, Azaleas. Birches, Dogwoods, Junipers, Lilacs, Maples, Mums, Oaks, Pines, and Roses. Do not use on Boston Ivy. Activate attached sprayer and apply 8 fl oz. cf this product to treat 1,333 sq. ft. or simply spray plants to point of spray runoff. Apply when insects or damage appear, for such insects as Bagworms, Elm Leaf Beetles, Gypsy Moths, Japanese Beetles, Lace Bugs, Leafminers, Leaf Rollers. Periodical Cicadas, Rose Slugs, Scale Insects, and Tent Caterpillars. Repeat as necessary for control, but not more than once a week.

Vegetable Garden Spray: Activate attached sprayer and apply 8 ozs. Of this product to treat 1,333 sq. ft. Spray when insects or damage appear. Do not apply within 3 days of harvest.

Beans (green, lima and snap): Japanese Bean Beetles, Mexican Bean Beetles, and Tarnished Plant Bugs.


Eggplants, Peppers, and Tomatoes: lace Bugs, Leafhoppers, and Tomato Fruit Worms.

Cucumbers, Melons, Pumpkins, and Squashes: Cucumber Beetles, Flea. Beetles, Melonworms, and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet, or in the presence of prolonged high humidity. Do not use on Watermelons in Floridations

SHAKE WELL BEFORE USING

This product is an extremely fine dispersion of carbaryl in water. In order to spray, activate the attached sprayer as directed and apply to thoroughly cover all plant surfaces. Occasionally shake or agitate sprayer unit during application. Avoid spray drift to adjoining food or forage crops in bloom.

4





5 95