



5%

Diazinon[®] Dust

Kills ants and lawn insects

Ready to use to kill insects; aphids, bagworms, imported cabbage worms, cutworms, mealybugs, millipedes, mites, and other listed insects on flowers, fruits, lawns, ornamentals, shrubs, and vegetables.



A - 5202B

Customers: Apply:
Aphids, bagworms, cutworms, mealybugs, millipedes, mites, and other listed insects on flowers, fruits, lawns, ornamentals, shrubs, and vegetables.

DO NOT APPLY TO ABOVE FOOD CROPS WITHIN 10 DAYS OF HARVEST

Customers: Apply:
Aphids, bagworms, cutworms, mealybugs, millipedes, mites, and other listed insects on flowers, fruits, lawns, ornamentals, shrubs, and vegetables.

DO NOT APPLY TO ABOVE FOOD CROPS WITHIN 10 DAYS OF HARVEST

Customers: Apply:
Aphids, bagworms, cutworms, mealybugs, millipedes, mites, and other listed insects on flowers, fruits, lawns, ornamentals, shrubs, and vegetables.

DO NOT APPLY TO ABOVE FOOD CROPS WITHIN 21 DAYS OF HARVEST

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid breathing of dust. May be absorbed through skin. Avoid contamination of food and feed. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Avoid storage near food and feed products.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
LLGIM, IL'INOIS 301,0
00-385

25/0674

Black Leaf®

SINCE 1879

ACCEPTED

OCT 30 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5 587-104


Kills ants and lawn insects

ACTIVE INGREDIENT:
O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl)
phosphorothioate **5%**
INERT INGREDIENTS: **95%**
Total **100%**

DIAZINON® is a Registered trade mark of Ciba-Geigy Corporation;
 U.S. Patent No. 2,754,243.

CAUTION
KEEP OUT OF
THE REACH OF CHILDREN
See other cautions on rear panel.

EPA REG. NO. 5887-104 AA
Net Weight 16 oz. (1 pound)



DIRECTIONS FOR USE OF BLACK LEAF 5% DIAZINON[®] DUST

DIRECTIONS FOR USE OF BLACK LEAF 5% DIAZINON* DUST
Please read entire label observing all cautions. Apply BLACK LEAF 5% DIAZINON DUST as soon as insects appear or their damage is noticeable. On soil or lawns, spread evenly. For use on foliage, dust evenly covering both upper and under sides of foliage. Repeat as often as necessary to maintain attractive appearance of garden. Refer to table for more detailed instructions and limitations. Where instructions are given for ounces, please figure that one ounce equals four tablespoonsful.

| USE | INSECTS CONTROLLED | APPLICATION INSTRUCTIONS |
|---|--|---|
| ORNAMENTALS | Aphids, bagworms, carnation shoot mites, clover mites, cyclamen mites, European red mites, flea beetles, holly bud moths, leafhoppers, Mimosa-webworms, scale crawlers, tent caterpillars, thrips, two-spotted mite, whiteflies. | Dust evenly, covering both upper and undersides of foliage. For best results, treat when air is still. Do not spray when temperature exceeds 90° F. Do not use on ferns, poinsettias, orchids, pines, and papaya. Since injury to plants may occur. |
| LAWNS | Ants, chiggers, fleas, leafhoppers, ticks | Spread 1 1/4 oz. per 125 sq. ft. area. Water lawn after application. Repeat as necessary. |
| | Clover mites, crickets, | Spread 4 oz. per 125 sq. ft. area. Water lawn after application. Repeat as necessary. |
| | Millipedes | Use 8 oz. per 125 sq. ft. area. Water lawn after application. |
| SOIL INSECTS | Cutworms, onion maggots, root maggots | Spread 2 oz. per 100 sq. ft. area. Work into the soil immediately. 2-3 inches. |
| | Mole crickets | Spread 1 oz. per 100 sq. ft. and work into the soil immediately. 1-2 inches. |
| VEGETABLES AND MELONS | Aphids, leafhoppers, mites, thrips | For foliar application, apply dust evenly on upper and lower surfaces. Repeat application as necessary. |
| Cantaloupe | | |
| Peas | Aphids, thrips | |
| Sweet corn | Aphids, corn earworm | |
| Tomatoes | Dipterous leaf miners, mites | |
| Watermelon | Aphids, leafhoppers, thrips | |
| DO NOT APPLY TO ABOVE FOOD CROPS WITHIN 3 DAYS OF HARVEST. | | |
| Broccoli | Imported cabbageworms | For foliar application, apply dust evenly on upper and lower surfaces. Repeat application as necessary. |
| Cabbage | Aphids | |
| Cauliflower | Aphids | |
| Celery | Aphids | |

DO NOT APPLY TO ABOVE FOOD CROPS WITHIN 3 DAYS OF HARVEST

| | | |
|-------------|-----------------------|---|
| Broccoli | Imported cabbageworms | For foliar application, apply dust evenly on upper and lower surfaces. Repeat application as necessary. |
| Cabbage | Aphids | |
| Cauliflower | Aphids | |
| Celery | Aphids | |