


EPA EST. NO. 5887-1L-1

Black Leaf[®]
SINCE 1879

Kills dandelions and weeds

ACTIVE INGREDIENTS:
 *Isooctylester of Silvex
 (2-(2,4,5-Trichlorophenoxy)
 propionic acid) 4.94%
 **Isooctylester of
 2,4-Dichlorophenoxy
 acetic acid 15.84%
INERT INGREDIENTS 79.12%
TOTAL 100.00%



*Equivalent to 3.49% Silvex
 (2-(2,4,5-Trichlorophenoxy)
 propionic acid) or 0.28
 pounds per gallon.
 **Equivalent to 18.87% 2,4-
 Dichloro-phenoxyacetic acid
 or 0.86 pounds per gallon.

CAUTION
KEEP OUT OF THE
REACH OF CHILDREN
 See rear panel for other cautions

EPA Reg. No. 5887-106 AA Net Contents 32 fl. oz. (1 quart)

ACCEPTED
 FEB 27
 Under the Federal
 Pesticide Act
 as amended
 registered under
 EPA Reg. No. 5887-106
 Comments

GENERAL DIRECTIONS FOR USE
BLACK LEAF WEED KILLER contains both 2,4-D and Silvex for control of a broad range of weeds infesting lawns.
BLACK LEAF WEED KILLER is effective for the control of cow, clover, dock, dog fennel, English daisy, Frenchweed, knotweed, radish, ground ivy, houndstooth, knot-grass, lamb's-ear, morning glory, chickweed (many), scarab, smartweed, oxalis, pennywort, poppygrass, pigweed, purslane, p. oak, Canada thistle, burdock, groundsel, purple top, ragwort, sheep sorrel, spotted spurge, wild radish, wild rose, wild yam, wild onion, wood sorrel, bindweed, and yarrow.
LAWN WEEDS: To kill dandelion, wild oxalis, pennywort, and similar weeds, use 8 ounces in 6 gallons of water in a spray or sprinkling can to 2,500 square feet of lawn. (For use 2 teaspoonsful in 2 quarts of water applied to 100 square feet.) Do not apply to areas underlain by roots of desirable trees.
 To kill weeds such as Canada thistle, bindweed, dock, and similar weeds, use 8 ounces in 6 gallons of water (or 2 teaspoonsful in 2 quarts of water) applied directly to plants, wetting them thoroughly. Repeat as needed.
 To kill undesirable plants such as poison ivy and poison oak, use 2 teaspoonsful in 2 quarts of water and drench plants when they are in leaf. Repeat in about 4 weeks if necessary. Certain plants such as poison ivy, poison oak, English daisy may require a second application.
IMPORTANT: Avoid drift of spray (coarse sprays are less desirable). Do not contaminate domestic or irrigation water used to clean sprayer near desirable plants or water used to wash. Do not store in direct contact with fertilizers, insecticides, or fungicides. Do not use on centipede grass, bermudagrass, or on lawns or turf where desirable bent grass is present. Newly seeded lawns may be treated after they have established the second time.
 To re-seed small areas in treated lawns, wait at least 3 weeks. A good rain or deep watering should occur during this period. If container is empty, immediately wash thoroughly and destroy. Use spray equipment for applying 2,4-D combinations only. Vapors from this product at very high temperatures may be harmful to susceptible plants growing nearby. Do not use spray equipment which has contained this product, for other purposes.
CAUTION: Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water; for eyes, get medical attention immediately.
NOTICE: Buyer assumes all risks of use, storage or handling. Use not in strict accordance with directions given herewith.


DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
 ELGIN, ILLINOIS 60120
 00-803

ack®
af

1879

dandelions and weeds

INGREDIENTS:
 Ester of Silvex (2,4,5-Trichlorophenoxyacetic acid) 4.94%
 Ester of Dichlorophenoxyacetic acid 15.84%
INGREDIENTS 79.12%
TOTAL 100.00%



*Equivalent to 3.46% Silvex (2-(2,4,5-Trichlorophenoxy) propionic acid) or 0.28 pounds per gallon.
 **Equivalent to 10.57% 2,4-Dichlorophenoxyacetic acid or 0.86 pounds per gallon.

ACCEPTED
 FEB 27
 Under the Federal Insecticide, Fungicide, and Rodenticide Act registered under EPA Reg. No. 5887766
 Comments

UTION
UT OF THE
OF CHILDREN
 See label for other cautions

g. No. 5887-106 AA Net Contents 32 fl. oz. (1 quart)

GENERAL DIRECTIONS FOR USE

BLACK LEAF WEED KILLER contains both 2,4-D and Silvex for effective control of a broad range of weeds infesting lawns.
BLACK LEAF WEED KILLER is effective for the control of dandelion, chickweed, clover, dock, dog leaflet, English daisy, Frenchweed, knotweed erect, knotweed prostrate, ground ivy, hawkweed, heat-ill, horsetail, lamb's-quarter, milkweed, morning glory, chickweed (most common-stiffwort); nuttall, oxalis, pennywort, peppercress, plantain, poison ivy, poison oak, Canada thistle, burdock, garden weed, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, white clover, wild carrot, wild radish, wild onion, wild rangel, wood sorrel, bindweed, and yellow.
LAWN WEEDS: To kill dandelion, wild onion, pennywort, plantain, heat-ill and similar weeds, use 8 ounces in 6 gallons of water and apply with sprayer or sprinkling can to 2,500 square feet of lawn. (For smaller areas use 2 teaspoonsful in 2 quarts of water applied to 100 square feet.)
 Do not apply to areas underlain by roots of desirable trees and shrubs.
 To kill weeds such as Canada thistle, bindweed, dock, and ragweed, use 8 ounces in 6 gallons of water (or 2 teaspoonsful in 2 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary.
 To kill undesirable plants such as poison ivy and poison oak, use 8 teaspoonsful in 2 quarts of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard to kill weeds such as poison ivy, poison oak, English daisy may require a fall application.
IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides. Do not use on centipede grass, St. Augustine or dichondra, nor on lawns or turf where desirable bent grass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time.
 To re-seed small areas in treated lawns, wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period. When container is empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.
 Vapors from this product at very high temperatures may cause injury to susceptible plants growing nearby. Do not use spray equipment which has contained this product, for other purposes.
CAUTION: Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.
CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with abundance of water; for eyes, get medical attention. Keep out of reach of children.
NOTICE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
 ELGIN, ILLINOIS 60120
 00-803

EPA EST. NO. 6887-11-1

Black Leaf[®]

SINCE 1879


Kills dandelions and weeds

ACTIVE INGREDIENTS:

*Isooctylester of Silvex (2-(2,4,5-Trichlorophenoxy) propionic acid)	4.94%
**Isooctylester of 2,4-Dichlorophenoxy acetic acid	15.94%
INERT INGREDIENTS ...	79.12%
TOTAL	100.00%

CAUTION
KEEP OUT OF THE
REACH OF CHILDREN

See rear panel for other cautions



*Equivalent to 3.49% Silvex
(2-(2,4,5-Trichlorophenoxy)
propionic acid) or 0.28
pounds per gallon.

**Equivalent to 10.57% 2,4-
Dichlorophenoxyacetic acid
or 0.88 pounds per gallon.

EPA Reg. No. 5887-106 AA Net Contents 32 fl. oz. (1 quart)

Black Leaf

SINCE 1871


Kills dandelions and weeds

ACTIVE INGREDIENTS


Isopropyl ester of Silvex	
2-(2,4,5-Trimethylphenoxy)	4.94%
2,4-Dichlorophenoxy	1.84%
NET CONTENTS	29.12%
TOTAL	

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
 See rear panel for other cautions

EPA Reg. No. 5887-106 AA Net Contents 32 fl. oz. (1 quart)



Black Leaf Weed Killer kills dandelions and weeds.



GENERAL DIRECTIONS FOR USE

BLACK LEAF WEED KILLER contains both 2,4-D and Silvex for effective control of a broad range of weeds infesting lawns.

BLACK LEAF WEED KILLER is effective for the control of dandelion, chicory, clover, dock, dog fennel, English daisy, Frenchweed, knotweed erect, knotweed prostrate, ground ivy, hawkweed, heal-all, henbit, lambsquarter, mallow, morning glory, chickweed (mouse ear), common-stitchwort, mustard, oxalis, pennywort, peppergrass, pigweed, plantain, poison ivy, poison oak, Canada thistle, burdock, jimson weed, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, wild astor, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, bindweed, and yarrow.

LAWN WEEDS: To kill dandelion, wild onion, pennywort, plantain, heal-all and similar weeds, use 8 ounces in 6 gallons of water and apply with sprayer or sprinkling can to 2,500 square feet of lawn. (For smaller areas use 2 teaspoonsful in 2 quarts of water applied to 100 square feet.)

Do not apply to areas underlaid by roots of desirable trees and shrubs.

To kill weeds such as Canada thistle, bindweed, dock, and ragweed, use 8 ounces in 6 gallons of water (or 2 teaspoonsful in 2 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary.

To kill undesirable plants such as poison ivy and poison oak, use 8 teaspoonsful in 2 quarts of water and spray plants when they are in flower.

Repeat application about 4 weeks later, if necessary, to control regrowth.

For information on the use of Black Leaf Weed Killer on other weeds, consult the label on the product.

Black Leaf Weed Killer is a registered trademark of the manufacturer.

© 1971 by the manufacturer. All rights reserved.