Do not use on household pets or humans. Do not permit children

or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

FISH AND WILDLIFE WARNING: This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. So net apply where runoff is likely to occur. This product is highly toxic to pollinating bees exposed to direct treatment or residues on cross. Protective information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate ornamental fishponds.

CAUTION: Do not re-use empty container. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. KEEP OUT OF SMOKE.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



Also controls sod webworms

Kills bagworms on evergreens, ornamentals, and shade trees, Also kills and controls sod webworms and billbugs in lawn grasses.



DIRECTIONS FOR USE

Step 1: Determine size of area to be sprayed for lawns or ornamentals.

Step 2: Use a clean sprayer. Carefully measure and mix the amount of bagworm spray and water as suggested in the directions.

Step 3: Spray as directed. Use coarse drenching sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

DO NOT USE IN THE HOME.

CAUTION: Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyels and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

Do not use on household pets or humans. Do not permit child:en or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

FISH AND WILDLIFE WARNING: This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. This product is highly toxic to pollinating bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate ornamental fishponds.

CAUTION: Do not re-use empty container. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning KEEP OUT OF SMOKE.

NOTICE. Buyer assumes all risks of use storage or handling of this material not in strict accordance with directions given herewith



ACCEPTE

DEC 1 5 1975

Under the Federal Insecticide Pungicide, and Redenticide a amended, for the posticide registered under EPA Reg. No. 5-557-//

Also controls sod wer wor

ACTIVE INGREDIENTS: By Wt 0.0-diethyl 0-(2-isopropyl-6-methyl 4-pyrimidinyl) phosphorothioate* 12.5% Aromatic Petroleum Derivative Solvents 72.6% INERT INGREDIENTS: 14.9%

Contains Diazinon. A registered trade mark of Ciba-Geigy Corp

CAUTION KEEP OUT OF THE REACH OF CHILDREN See side panel for other cautions



EPA REG. NO. 5887 Net Contents 16 fl. o and mix the amount the directions.

Ing sprays for trees
Flush sprayer with

ME.

orbed through skin.

with skin, eyes and

void contamination

not permit child ten intil the spray has stic animals.

AR HEAT OR OPEN

is toxic to fish, birds areas may be killed apply when weather of apply where runoff to pollinating bees ops. Protective infore Agricultural Extensified on this label.

uipment or disposal shponds.

inse equipment and ng in non-croplands d be disposed of by astes or by burning.

ge or handling of this ons given herewith



ACCEPTED

DEC 1 5 1975

Under the Federal Insections.
Pungislde, and Redenticide Act
as amended, for the posticide
revistared under
EFA Reg. No. 5-557-107

Also controls sod webworms

ACTIVE INGREDIENTS:

0.0-diethyl 0-(2 isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 12.5%

Aromatic Petroleum Derivative Solvents 72.6%

INERT INGREDIENTS: 14.9%

TOTAL 100.0%

Contains Diazinon. A registered trade mark of Ciba-Geigy Corp

CAUTION
KEEP OUT OF THE
REACH OF CHILDREN
See side panel for other cautions



EPA REG. NO. 5887-109 AA Net Contents 16 fl. oz. (1 pint)

DIRECTIONS FOR USE

BAGWORMS: For protection from bagworms on ornamentals such as arborvitae, douglas fir, juniper, locust, maple,
ook, pine and spruce.

Use 2 fluid ounces (4 tablespoonsful) in 3 gallons of water and apply as a thorough foliar spray. Wet leaves and branches to the dripping point and try to penetrate dense foliage in order to accomplish adequate protection. Upper and lower leaf surfaces should be treated. For best results apply when air is still.

SOD WEBWORMS: To protect lawns from sod webworms use 2 dunces (4 tablespoonsful) in 3 gallons of water to cover 125 sq. ft. area. 1 pint of BLACK LEAF BAGWORM SPRAY can be used in 24 gallons of water and will give protection on up to 1000 sq. rt. of lawn area. Thoroughly wet down the grass before applying.

BILL BUGS: For bill bug control apply 2 ounces in 3 gallons of water to cover 125 sq. ft. area or 16 ounces in 24 gallons of water to cover 1000 sq. ft. area. Water lawn thoroughly after treatment

Do not use on certain ferns, poinsettia hibiscus, papaya, pilea and gardenia and orchids since injury may occur

The American Assistance of the control of the second

90.488

Do not use on household pels or humans. Do not permit cl. ii. en or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

DO NOT USE. POUR. SPILL OR STORE NEAR HEAT OR OPEN FLAME.


FISH AND WILDLIFE WARNING: This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. This product is highly toxic to pollinating bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate ornamental fishponds.

CAUTION: Do not re-use empty container. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. KEEP OUT OF SMOKE.

NOTICE Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here with.

Also controls sod webworms



SOD WEBWORMS: To protect lawns from sod webworms use 2 ounces (4 tablespoonsful) in 3 gallons of water to cover 125 sq. ft. area. 1 pint of BLACK LEAF BAGWORM SPRAY can be used in 24 gallons of water and will give protection on up to 1000 sq. ft. of lawn area. Thoroughly wet down the grass before applying.

BILL BUGS: For bill bug control apply 2 ounces in 3 gallons of water to cover 125 sq. ft. area or 16 ounces in 24 gallons of water to cover 1000 sq. ft. area. Water lawn thoroughly after treatment.

Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia and orchids since injury may occur.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY

00-388

DIRECTIONS FOR USE

Step 1: Determine size of area to be sprayed for lawns or ornamentals.

Step 2: Use a clean sprayer. Carefully measure and mix the amount of bagworm spray and water as suggested in the directions.
Step 3: Spray as directed. Use coarse drenching sprays for trees

Step 3: Spray as directed. Use coarse drenching sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

DO NOT USE IN THE HOME.

CAUTION: Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

Do not use on household pets or humans. Do not permit of IId and pets to go deto the sprayed grass until the spray has ampletely great Keep out of reach of growestic aremais.

NO NOT USE PORT SPILE OF STORE NEAR HEAT OF THEN STAME

The second of the Bridge feed reading of treated reads may be killed as a control with the Bridge feed reading treated reads may be killed as a control records for manager pends. Commany pends represented apply where tanether here, the permitter marginated is a comparable apply where tanether here, and the permitter of the pends of the pends

The state of the s



ACCEPTED

DEC 1 5 19/5

Under the Federal Insecticite, Pungicide, and Redenticide Act, on amended, for the posticide registered under EPA Reg. No. 575 § 7 -167

Also controls sod webworms

ACTIVE INTEREDIENT

O serby Geolder production of a company of the production of the company of

CAUTION
KEEP OUT OF THE
REACH OF CHILDREN
See side panel for other cautions



EPA REG. NO. 5887-109 AA Net Contents 16 fl. oz. (1 pint)

DIRECTIONS FOR USE

BAGWORMS: For protection from bagworms on ornamentals such as arborwitae, douglas fir, juniper, locust, maple,
co oak, pine and spruce.

Use 2 fluid ounces (4 tablespoonsful) in 3 gallons of water and apply as a thorough foliar epray. Wet leaves and branches to the dripping point and try to penetrate dense foliage in order to accomplish adequate protection. Upper and lower leaf surfaces should be treated. For best results apply when air is still.

SOD WEBWORMS To protect lawns to miscod webworms use 2 cumer 2 concerns at 25 sq. ft. area. 1 port of Brack. LEAF BAGWORM SPRAY are to use 1 on 24 quilibrium of wetter and with question 24 quilibrium of the transfer of the area. The concerns of the transfer of the area of the area. The concerns of the area of the area.