

Reg. # 5887-120

KA-17

182

Reviewed. Registrant claims to be in accordance with
regulation of Federal Insecticide, Fungicide and Rodenticide Act, 28 CFR 2.12, Additive to Inert Change in Status.

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Keep out of reach of children. Avoid breathing spray or contact with eyes or skin. Wash thoroughly after use. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not use near fishbowls or aquaria. Do not apply directly to water.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label and apply as directed. Shake well before using. This insecticide may be used outdoors in gardens or in greenhouses throughout the growing season or as a pre harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

For best results, use throughout the growing season to prevent buildup of insect populations. Hold can in an upright position while spraying. Spray plants from a distance of 1 1/2 to 2 feet using short bursts (one second). To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of the leaves. Contact insects directly whenever possible. Repeat applications at 7 to 10 day intervals, after heavy rains, or as reinfestation occurs.

USE ON: asparagus, beans, broccoll, cabbage, brussels sprouts, cauliflower, celery, cranberries, cucumbers, melons, squash, eggplants, lettuce, mustard greens, kale, collards, turnips, peppers, potatoes, radishes, spinach, sweet corn and tomatoes.

TO CONTROL: asparagus beetles, aphids, leafhoppers, whiteflies, Mexican bean beetles, 12 spotted cucumber beetles, cabbage loopers, cross striped cabbageworms, diamondback moth caterpillar, flea beetles, harlequin bugs, imported cabbageworms, stink bugs, green peach aphids, leafllers, fireworms, leafhoppers, blister beetles, Colorado potato beetles, webworms, Japanese beetles.

STORAGE: To be placed in areas inaccessible to children. Also, see above precautionary statements. **CONTAINER DISPOSAL:** Replace cap and discard container in trash. Do not incinerate or puncture.

NOTICE: Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given herein.

DISTRIBUTED BY
BLACK LEAF PRODUCTS COMPANY
ELGIN, ILLINOIS 60120
© 1977



Tomato & Vegetable Spray


For healthier vegetables

For use on tomatoes, potatoes, peppers, melons, eggplant, squash and sweet corn up to day of harvest.

Contains Natural Pyrethrum

ACTIVE INGREDIENTS:	
Pyrethrins	0.050%
Piperonyl Butoxide	
Technical	0.500%
Petroleum Distillate	0.825%
INERT INGREDIENTS:	98.625%
TOTAL	100.000%

Equivalent to min. 0.40% (butyl carbetyl (6 propylpiperonyl) ether and 0.10% other related compounds.



KEEP OUT OF REACH OF CHILDREN
CAUTION

See rear panel for additional precautionary statements.

EPA EST. NO. 334-IL-2
EPA REG. NO. 5887-120
Net Weight 12 oz.

ATTACHMENT I

A. ACTIVE INGREDIENTS

Pyrethrins	0.050%
* Piperonyl Butoxide Technical	0.500%
Inert Ingredients	99.450%
	<hr/>
Total	100.00%

*Equivalent to min. 0.40% (butylcarbityl)
(6 propylpiperonyl) ether and 0.10% other
related compounds.

B. Black Leaf Products
A Division of Wilbur-Ellis Company
Buckner, Kentucky 40010