

PM 19

5887-162

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

page 195  
5514680  
200  
17

FEB 24 1997

Stacy A. Schmidt  
SureCo  
10012 N. Dale Mabry Hwy.  
Suite 221  
Tampa, FL 33618

Subject: Black Leaf Liquid Fruit Tree Spray with Fungicide  
EPA Registration No. 5887-162  
Amendment dated November 14, 1996  
Response to Agency letter dated 10/1/96

Dear Ms. Schmidt:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:

- a) Move the two paragraphs located directly beneath the User Safety Requirements box ("Apply when air is calm to avoid.....applications more than once a week.") as required in the Agency letter dated 10/1/96 to the DIRECTIONS FOR USE section.
- b) Because this product is a liquid spray, you must change the sentence "Allow dust to settle in treated areas before reentering" to "Allow spray to dry in treated areas before reentering."
- c) Correct the typographical error in the sentence "Spray thoroughly to wet upper sand lower leaf...." Change "sand" to "and."
- d) Change "5-7" to "7" in the sentence "When insects are seen, shorten spray intervals to 5 7 days."

CONCURRENCES

SYMBOL	SURNAME	DATE						

F. Edwards A-1 5887/162.1 2/13/97

2 9 5

-2-

2: Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

/s/

Dennis H. Edwards, Jr.  
Product Manager (19)  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

Enclosure

3 9 5

# BLACK LEAF LIQUID FRUIT TREE SPRAY WITH FUNGICIDE

Protects fruit and trees 5 Ways!

A complete, water-sprayable concentrate containing Insecticide, Fungicide, Aphicide, Miticide, and Scalicide for broad-spectrum kill that saves the fruit for you.

Active Ingredients:

*Mehtoxychlor [1,1,1-trichloro-2,2-bis (p-methoxyphenyl)ethane] .....	12.00%
Malathion [0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate ...	6.00%
Carbaryl (1-naphthyl N-methylcarbamate) .....	0.30%
Captan [N-trichloromethyl thio-4-cyclohexene-1, 2-dicarboximide .....	11.74%
Related Derivatives .....	0.26%
Inert Ingredients: .....	<u>69.70%</u>
TOTAL:	100.00%

Contains xylene range aromatic solvent.

\*Equivalent to 10.56% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.44% of other isomers and related compounds

EPA Reg. No. 5887-162

EPA Est. No. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See Back Panel For Additional Precautionary Statements

### STATEMENT OF PRACTICAL TREATMENT

**If In Eyes:** Flush eyes immediately with water. Remove any contact lenses, then hold eyelids open and flush eyes with plenty of water. Call a physician.

**If Swallowed:** Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

**If Inhaled:** If ill or unconscious, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**If On Skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

Net Contents: \_\_\_\_\_


**ACCEPTED  
with COMMENTS  
in EPA Letter Dated**

**FEB 24 1997**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

5887-162

Manufactured for:  
Black Leaf Products  
Louisville, KY 40222



495

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin or clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic invertebrates, and toxic to fish and aquatic life stages of amphibians. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

The complete, easy-to-use concentrate that mixes with water instantly; no plugged nozzles; no messy powders to handle, measure or mix; no premixing or straining necessary. Designed especially for home orchards only to protect Apples, Cherries, Grapes and Peaches from the ravages of insects and diseases listed below.

### USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

*Spray to dry*

For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper sand lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**INSECTS AND DISEASES CONTROLLED APPLES:** Aphids (Woolly Apple, Green Apple, and Rosy Apple), Apple Maggots, Bud Moths, Coding Moths, Eastern Tent Caterpillars, Forbes Scales, Japanese Beetles, Leafhoppers, Mealybugs, Mites (Clover, European Red, Two-Spotted Spider, and Willamette), Plum Curculios, and Red-Banded Leaf Rollers. Bitterrot, Black Rot (Frogeye), Potryosphaeria Rot, Botrytis (Blossom- End) Rot, Brooks Fruit Spot, Fly Speck, Scab and Sooty Blotch.

**CHERRIES:** Black Cherry Aphids, Cherry Fruit Flies, Cherry Fruit Worms, Fruit Tree Leaf Rollers, Japanese Beetles, and Plum Curculios. Black Rot, Botrytis (Bunch) Rot, and Downy Mildew.

**GRAPES:** Flea Beetles, Grape Berry Moths, Japanese Beetles, Leafhoppers, Mealybugs, and Spider Mites. Black Rot, Downy Mildew (berries only), Botrytis rot.

595

**PEACHES:** Aphids (Black Cherry, Black Peach, European Red, Green Peach, and Rusty Plum), Cottony Peach Scales, Japanese Beetles, Oriental Fruit Moths, Plum Curculios, Spider Mites, and Terrapin Scales. Brown Rot (Blossom Blight), Coryneum Blight (Peach Blight), Shothole, Scab.

**SHAKE PRODUCT THOROUGHLY BEFORE USING.** Contains extremely fine particles which settle upon standing, and require reblending by agitation. Choose a cool, calm period, preferably early morning or evening, when trees will not be subjected to hot, direct rays of the sun. Do not apply if rain is expected before leaf surfaces dry. It is easier to prevent insects and scale damage than to cure them, therefore a preventative 10-14 day spray schedule is recommended. When insects are seen, shorten spray intervals to 5-7 days. Apply spray to leaves, fruit, branches, trunk, and base of tree until lightly dripping. Spray 1-2 pints per small/dwarf tree, 1-2 quarts per medium size tree, and 1-3 gallons per large/mature tree. Shake sprayer occasionally (or agitate) to keep spray particles in suspension during application. Follow spray schedule given in spray chart, using tablespoonfuls (15 ml) per gallon (3.8 L) of water. Observe precautionary statements and cautions for fruit to be treated.

**SPRAY CHART**

Add amount indicated below to 1 gal. water. (1 tablespoonful (T.) = 1/2 fl oz = 15 ml)

TREE STAGE	APPLE*	CHERRY	GRAPE*	PEACH
Dormant	Use a dormant oil spray up to 7 days before green tip application. Start and repeat spraying at 10 day intervals until bloom.			
Green tip	1 1/2 T.	1 1/2 T.	1 1/2 T.	1 1/2 T.
Pre-bloom	2 1/2 T.	1 1/2 T.	2 1/2 T.	1 1/2 T.
Full pink	2 1/2 T.	2 1/2 T.	2 1/2 T.	2 1/2 T.
Bloom	Don't spray pollinators. Fungicide alone may be applied for disease. Repeat at 14 day intervals, stop at days before harvest shown below.			
Petal fall	2 1/2 T.	2 1/2 T.	2 1/2 T.	2 1/2 T.
1st cover	2 1/2 T.	2 1/2 T.	2 1/2 T.	2 1/2 T.
2nd to Harvest	2 1/2 T.	2 1/2 T.	2 1/2 T.	2 1/2 T.
	Stop spraying 14 days before harvest of Apples, Cherries or Grapes; 21 days before for Peaches.			
Post harvest	1 1/2 T.	1 1/2 T.	1 1/2 T.	2 1/2 T.

\*NOTE: Malathion may injure foliage on some varieties of Apples, and Grapes.

**Definitions of terms**

**Dormant** - absence of growth in fall, winter, or spring. **Green tip** - when branch tips show green tissue. **Pre-bloom** - when blossom buds begin to show color. **Full pink** - when blossom buds show full color before opening. **Bloom** - when blossoms are open. **Petal fall** - when last petals have fallen from blossom. **1st cover** - 7 to 10 days after petal fall. **2nd cover** - to cut-off date before harvest. **Post harvest** - to leaf drop.

**STORAGE AND DISPOSAL**

**Storage:** Keep in upright, tightly closed original container in a cool, dry, locked area away from children and domestic pets. **Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container Disposal:** Do not reuse empty container. Rinse original container thoroughly, then securely wrap in several layers of newspaper and discard in trash.