

BLACK LEAF LAWN & GARDEN FUNGICIDE

A Broad Spectrum fungicide for use on turf, grass, dichondra, vegetables and
ornamentals.

ACTIVE INGREDIENT:

Chlorothalonil (tetrachloroisophthalonitrile)* 12.5%

INERT INGREDIENTS: 87.5%

TOTAL: 100.0%

* Daconil 2887 covered under U.S. Patent No. 3,290,353 and No. 3,331,735
This product contains 1.14 pound Chlorothalonil per gallon.

EPA Reg. No. 5887-176

EPA Est. No. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN
WARNING**

See Additional Precautionary Statements on Back Panel

Net Contents: 1 Quart (946 ml)

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

WARNING: Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Do not breathe spray mist. Can be irritating to eyes or skin. Skin sensitizer. Wash thoroughly with soap and warm water after handling.

NOTE: Black Leaf Lawn and Garden Fungicide may produce temporary allergic side effects characterized by redness or rash on exposed skin area. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroid. There is no evidence of after effects of permanent injury.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until fluid is clear. Call physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air. Apply respiration if indicated.

If On Skin: Wash thoroughly with soap and warm water after handling.

If In Eyes: Flush with plenty of water immediately for at least 15 minutes. Call a physician.

Received 10/11/97

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Keep children and pets away from treated area until it is completely dry.

SHAKE WELL BEFORE USE

Read entire label and use as directed. Apply only to areas specified on this label. Black Leaf Lawn and Garden Fungicide is highly effective and widely recommended for the control of a number of important lawn and ornamental commonly occurring diseases. Do not combine this product with pesticides, or fertilizers unless your prior use has shown the combinations physically compatible and non-injurious under your conditions of use.

Note To User: Do not graze treated areas. Do not feed clippings to livestock.

Note To User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note To User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

GRASS LAWNS: For control of Brown Patch, Dollar Spot, Leaf Spot, "Blight, Fading out, Melting Out", Gray Leaf Spot, "Stem Rust on Bluegrass" and Copper Spot. Mix 1 pint in 5-15 gallons of water and apply to 500 sq. ft. of lawn. Start spray application in spring before disease appears and repeat at 7-14 day intervals throughout the season. Make application on 7 day schedule during periods of wet weather or severe disease condition. Do not mow or water after treatment until lawn is thoroughly dry.

DICHONDRA LAWNS IN CALIFORNIA ONLY: To cure Alternaria Leaf Spot use 1 pint of material in 5 to 10 gallons per 500 sq. ft. Spray on 7-14 day intervals of application. Under severe disease conditions spray on a 7 day schedule and use higher rates. Use lower rates when disease conditions are light to moderate.

ORNAMENTALS: Black Leaf Lawn & Garden Fungicide is recommended for the control of a number of diseases on ornamentals. Apply at first sign of disease and repeat at 7 to 14 day intervals throughout the season. Applications should be thorough to obtain adequate coverage of the foliage.

CHRYSANTHEMUMS: Ascochyta Ray Blight and gray mold use 9 tablespoonfuls per gallon of water.

GERANIUMS: Botrytis Blight use 9 tablespoonfuls per gallon of water.

IRIS: Botrytis Blight and leaf spot use 9 tablespoonfuls per gallon of water.

ROSES: Black spot and blossom blight use 6 tablespoonfuls per gallon of water.

ZINNIAS: Powdery mildew use 9 tablespoonfuls per gallon of water.

PHILODENDRON: For control of Phytophthora Blight and Dactylaria leaf spot use 9 tablespoonfuls per gallon of water. Apply at 7 day intervals.

VEGETABLES:

Broccoli, Brussels Sprouts, Cabbage, Cauliflower. For *Alternaria* leaf spot (Southeastern states only) and downy mildew apply after transplant and when conditions favor disease development. Mix 3 tablespoonfuls in 5-15 gallons of water and apply to 500 sq. ft. Repeat at 7-10 day intervals or a necessary to maintain control.

MELONS (Cantaloupe, Honeydew, Muskmelon): For Anthracnose, downy mildew apply when conditions favor disease development. Use 6 tablespoonfuls in 5-15 gallons of water per 500 sq. ft. Repeat at 7 day intervals.

PUMPKIN, SQUASH (Summer and Winter), WATERMELONS: For Anthracnose, downy mildew apply when conditions favor disease development. Use 6 tablespoonfuls in 5-15 gallons of water and apply to 500 sq. ft. of garden area. Repeat at 7 day intervals.

TOMATOES: For Anthracnose, early and late blights, gray leaf spot, gray leaf mold use 4 tablespoonfuls in 5-15 gallons of water per 500 sq. ft. Begin when disease threatens and repeat at 7-10 day intervals.

STORAGE AND DISPOSAL

Store in a dry place. Do not contaminate water, food, or feed by storage and disposal. Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

WARRANTY AND LIMITATIONS OF DAMAGES: Seller warrants that this material conforms to its chemical description and is reasonable fit for the purpose stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions.

Distributed by:
Black Leaf Products
A Div. Of SureCo, Inc.
Louisville, KY 40222

