

ACCEPTED

OCT 16 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 58919 SUBJECT
TO ATTACHED COMMENTS.

SANI DET SANITIZER is recommended for use on food processing equipment and utensils, including dairy and brewery equipment and utensils. It is also recommended for use on dishes and glasses.

Directions for Use:

Prepare a sanitizing solution by dissolving 4 1/2 oz. or 8 1/2 oz. **SANI DET SANITIZER** per 10 gallons of water. This provides a solution containing 100 ppm or 200 ppm of available chlorine respectively. Clear and rinse items prior to sanitizing. Then place glasses, dishes, equipment or utensils in solution for 2 to 5 minutes. Remove, rinse with potable water and dry. For flat surfaces clean to remove gross filth and heavy soil. Then spread the sanitizing solution and allow to stand 2 to 5 minutes and then rinse with potable water. The same solution may be used in feed tank in immersion or spray type machines providing 2 to 5 minutes contact time for sanitizing dishes, glasses, food processing equipment and utensils. Clean and rinse items prior to sanitizing. After sanitizing step, rinse well with potable water and dry. The items are then ready for use.

The sanitizing solution should be used promptly after mixing. If the solution is allowed to stand or is reused and tests show the level of chlorine has not been maintained, additional **SANI DET SANITIZER** should be added, to maintain a minimum concentration of 100 ppm available chlorine.

SANI DET SANITIZER

ACTIVE INGREDIENTS:

Sodium dichloro-s-triazinetrione dihydrate
(provides 3.2% available chlorine)

Sodium dodecylbenzene sulfonate

Sodium metasilicate

5.60%

4.00%

5.00%

INERT INGREDIENTS:

85.40%

WARNING! KEEP OUT OF THE REACH OF CHILDREN.
See right side panel for precautionary and first aid statements.

M. HOOD CHEMICAL CORPORATION
PORTLAND, OREGON 97210



WARNING!

Directions for Storage:

Store in a cool, dry place. Contamination may result from chlorine containing gas. Do not mix with other chemicals. Do not use for may cause fire. Do not liberate of harmful gas. If possible isolate container. Flood with ventilated air. Flood with water if necessary.

Do not reuse empty container. Thoroughly wash clean. This product is toxic to fish. Do not be discharged where it will pollute ponds or public water. Do not use for cleaning of equipment or this product only as specified.

KEEP OUT OF THE REACH OF CHILDREN. May cause irritation if swallowed. May cause irritation in eyes, on skin, or on clothing.

FIRST AID!

If swallowed: feed with water or olive oil or cooking oil. Do not induce vomiting. If in contact with skin or eyes with fresh water. For eyes get medical attention. Rinse thoroughly with water.

E.P.A. Establishment No. 58919

E.P.A. Reg. No. 5891-19

Net Contents

Manufactured by
M. Hood Chemical Corporation
Portland, Oregon

SANIDET SANITIZER

ACTIVE INGREDIENTS

Sodium dichloro s triazinetrione dihydrate
(provides 3.2% available chlorine) 5.60%
Sodium dodecylbenzene sulfonate 4.00%
Sodium metasilicate 5.00%

INERT INGREDIENTS

85.40%

WARNING! KEEP OUT OF THE REACH OF CHILDREN
See right side panel for precautionary and first aid statements.

MT. HOOD CHEMICAL CORPORATION
PORTLAND, OREGON 97210



WARNING!

Directions for Storage & Handling:

Store in a cool, dry, well ventilated area. Moisture contamination may liberate irritating and hazardous chlorine containing gases. Oxidizing material. Do not mix with other chemicals. Contact with organic matter may cause fire. Decomposes at 350-475° F with liberation of harmful gases. In case of decomposition, if possible isolate container in open air or well ventilated area. Flood with large volume of water if necessary.

Do not reuse empty container. Rinse empty container thoroughly with clean water before discarding. This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. May cause eye damage. Do not get in eyes, on skin, or on clothing. Do not breathe dust.

FIRST AID!

If swallowed: feed gruel or cooked cereal followed by olive oil or cooking oil. Call physician. In case of personal contact: remove excess material and flush skin or eyes with fresh water for at least 15 minutes. For eyes get medical attention. Rinse empty container thoroughly with water and discard it.

E.P.A. Establishment No. 5891-OR-1

E.P.A. Reg. No. 5891-19

Net Contents

Manufactured by
Mt. Hood Chemical Corporation
Portland, Oregon 97210

Use on
sanitizing
It is also

1/4 oz. or
gallons of
100 ppm
Clean
glasses,
2 to 5
and dry.
no heavy
allow to
potable
tank in
2 to 5
glasses,
can and
ing step,
ems are

ly after
reused
t been
should
tion of

ACCEPTED

OCT 16 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. **589719** SUBJECT
TO ATTACHED COMMENTS.

SANI DET SANITIZER is recommended for use on food processing equipment and utensils, including dairy and brewery equipment and utensils. It is also recommended for use on dishes and glasses.

Directions for Use:

Prepare a sanitizing solution by dissolving 4 oz. or 8 oz. of SANI DET SANITIZER per 10 gallons of water. This provides a solution containing 100 ppm or 200 ppm of available chlorine respectively. Clean and rinse items prior to sanitizing. Then place glasses, dishes, equipment or utensils in solution for 2 to 5 minutes. Remove, rinse with potable water and dry. For flat surfaces clean to remove gross dirt and heavy soil. Then spread the sanitizing solution and allow to stand 2 to 5 minutes and then rinse with potable water. The same solution may be used in feed tank in immersion or spray type machines providing 2 to 5 minutes contact time for sanitizing dishes, glasses, food processing equipment and utensils. Clean and rinse items prior to sanitizing. After sanitizing step, rinse well with potable water and dry. The items are then ready for use.

The sanitizing solution should be used promptly after mixing. If the solution is allowed to stand for extended periods above the level of chlorine may not be maintained. Additional SANI DET SANITIZER should be added if the chlorine level is low. For more information contact your distributor or write to:

SANI SANITIZER

ACTIVE INGREDIENTS:
Sodium dichloro-s-triazinetriene
(provides 3.2% available chlorine)
Sodium dodecylbenzene sulfonate
Sodium metasilicate

INERT INGREDIENTS:

WARNING! KEEP OUT OF THE REACH OF CHILDREN
See right side panel for precautions

MT. HOOD CHEMICAL
PORTLAND, OREGON

