

## CAUTION

Read entire label. Use strictly in accordance with cautions and directions, and with applicable state and federal regulations.

Keep out of reach of children. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water. Fish may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not contaminate water by cleaning of equipment or disposal of wastes.

Keep pesticide in original containers. Burn bag immediately when empty. Stay out of smoke. Apply thoroughly. Do not apply to food crops within 3 days of harvest unless otherwise specified. This product is formulated for application by ground or air equipment.

### DIRECTIONS FOR USE

**ALFALFA:** Armyworms - 25 lbs. per acre. Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittlebugs, Vetch Bruchids - 30 lbs. per acre. Spotted Alfalfa Aphids - Ground - 12 to 15 lbs. per acre or Air - 12 to 20 lbs. per acre. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the field or are not hanging on outside of hives. May be applied on day of harvest or grazing.

**APPLES:** Red Spider Mites (such as European Red, Two-Spotted, Pacific, Willamette, Clover) (2 or more applications as needed). Green Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphids (1 or more applications as needed) - 30 to 40 lbs. per acre. This product may injure foliage on some varieties of apples.

**APRICOTS:** Spider Mites, Brown Apricot Scale (Crawlers), Orange Tortrix - 30 lbs. per acre. Do not apply within 7 days of harvest.

**ASPARAGUS:** Thrips - 20 to 25 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

**BEANS:** Mexican Bean Beetle, Japanese Beetles, Cucumber Beetles, Leafhopper, Spider Mites, Aphids - 30 to 35 lbs. per acre. Two or more applications as needed. Do not apply within 1 day of harvest.

**BLEUBERRIES:** Japanese Beetles - 20 lbs. per acre. Apply when berries turn blue and at 10-day intervals while infestation persists. Plum Curculio, Sharp-nosed Leafhopper - 32 to 40 lbs. per acre. Do not apply within 7 days of harvest.

**BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, KALE, MUSTARD GREENS, TURNIPS:** Aphids, Imported Cabbageworm, Cabbage Looper, Diamond-back Moth, Flea Beetles - 30 to 35 lbs. per acre. Repeat as necessary. Do not apply to Cauliflower, Brussels Sprouts, Cabbage, Mustard Greens and Kale within 7 days of harvest and to Broccoli and Turnips within 3 days of harvest.

**CARROTS:** Leafhoppers - 30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

**COTTON:** Boll Weevil, Aphid, Leafworm and Spider Mites - 20 to 30 lbs. per acre. Do not apply within 7 days of hand picking. The time between applications should not exceed 5 to 7 days. If applied by air, wind should be very light. Either morning or evening applications are recommended. Emphasis should be on complete coverage. Consult state agricultural extension service for timing of applications.

**CRANBERRIES:** Leafhoppers, Black-Headed Fireworms - Ground Machine - 30 lbs. per acre. Make thorough coverage.

**CUCUMBERS, SQUASH, MELONS:** Aphids, Spider Mites, Pickleworms, Squash Vine Borer - 30 to 35 lbs. per acre. CANTALOUPE - Leafhoppers - 30 to 35 lbs. per acre as necessary. Do not apply unless plants are dry nor within 1 day of harvest.

**DATES:** Nitidulid Beetles - Apply to cover each cluster thoroughly using 40 lbs. per acre. Make no application on dates within 7 days of the first picking.

**GRAPES:** Leafhoppers, Spider Mites - 25 to 30 lbs. per acre. Leafhoppers - One or more applications as needed. Spider Mites - Two or more applications as needed.

**LETTUCE:** Aphids, Mites, Cabbage Looper, Leafhoppers - 25 to 30 lbs. per acre. Do not apply to Head Lettuce within 7 days of harvest and to Leaf Lettuce within 14 days.

**MOSQUITOES:** Apply 10 to 15 lbs. per acre with a power duster. Avoid drift where susceptible fish and wildlife may be endangered.

**ONIONS (SEED ONLY):** Onion Thrips - 35 to 40 lbs. per acre.

# HELENA BRAND

## 5% CYTHON® DUST

THE PREMIUM GRADE MALATHION®

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL.**

ACTIVE INGREDIENT:	Malathion*	By Wt.	5%
INERT INGREDIENTS:			95%
	Total		100%

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

©Trademark of American Cyanamid Company.

MANUFACTURED BY  
**HELENA CHEMICAL COMPANY**

MEMPHIS, TENNESSEE

*Use Specific*  
**ORNAMENTALS:** Aphids, Red Spider Mites, Whitlily, Thrips, Japanese Beetle Adult, Four-lined Leaf Bug, Boxwood, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper - 25 to 30 lbs. per acre. Repeat as necessary. Do not use on Boston, Maidenhair and Pteris Ferns, and some species of Crassula.

**PEARS:** Pear Psylla, Aphids, Woolly Aphids - 30 to 40 lbs. per acre. One or more applications as needed. Spider Mites - Same directions listed under Apples. Do not apply within 1 day of harvest.

**PEAS:** Pea Aphid - 25 lbs. per acre as needed. Do not apply within 3 days of harvest or 7 days if forage is to be fed to livestock.

**PEPPERMINT, SPEARMINT:** Flea Beetle Adults - 20 lbs. per acre. Do not apply within 7 days of cutting.

**PEPPERS:** Pepper Maggots - 30 lbs. per acre.

**PUMPKINS:** Squash Vine Borer - 35 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply unless plants are dry.

**SPINACH:** Aphids - 25 to 30 lbs. per acre. Do not apply within 7 days of harvest.

**RASPBERRIES:** Aphids, Rose Scale - 35 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

**STRAWBERRIES:** Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Potato Leafhoppers, Strawberry Leafrollers, Whiteflies, Strawberry Root Weevil, Field Crickets - 40 lbs. per acre. Repeat as necessary. Strawberry Root Weevils - Apply to soil surface before planting and work into top 6 to 8 inches of soil.

**TOMACCO:** Aphids - 20 to 30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

**TOMATOES:** Aphids, Spider Mites, Tomato Russet Mite, Drasophila - 35 to 40 lbs. per acre. Do not apply within 1 day of harvest. **ANILLPEDES, SPRINGTAILS, SOWBUGS** In and Around Greenhouses, Around Homes, In Yards and Ornamental Gardens - 1 lb. per 150 sq. ft. (10 x 15') of soil surface or where insects congregate. Repeat at 7 to 10-day intervals as needed.

**FLEAS:** In Dog and Cat Kennels, In Pens, Yards, Lawns and under Houses - Use 1 to 2 lbs. per 1,000 sq. ft. Use a hand duster. Remove manure or debris before dusting.

**HORNFLIES:** Beef and Dairy Cattle - Apply 3 tablespoonfuls per animal. Do not apply more than 1.5 ounces per Dairy cow. Apply on back and neck of animals. Repeat at 10- to 14-day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not use sprays or dip dairy animals in malathion. Do not contaminate water troughs.

**LICE:** Swine, Beef Cattle - Dust animals thoroughly. Repeat in 10 days and as needed. For pigs less than 1 month of age, use 1/4 to 1/2 tablespoonful per animal. One treatment may be sufficient. Repeat only as needed. Do not treat calves under 1 month old. Avoid contamination of feed, food containers and watering troughs. Do not apply to dairy animals. Swine - Also treat pens thoroughly.

**BEEF AND NONMILKING CATTLE:** Hornflies, Lice Reduction of Fecundity (Dust-Bag Self Applicators) - Arrange dust-bag self applicators stacked with 10 to 20 lbs. of 5% Cythion Dust at mid-shoulder height near salt boxes or over fenced-in entrances to water troughs, ponds or loafing areas. Mandatory daily use will give the best results. Restock bags as frequently as necessary. Avoid contamination of salt, feed or water. Do not make accessible to lactating dairy animals or nonlactating animals within 2 weeks of freshening.

**POULTRY:** (Direct Application to Chickens, Geese, Ducks, Turkeys) Northern Fowl Mite, Poultry Lice, Chicken Red Mite (as a supplement to premise treatments for chicken red mite) - Dust individual birds thoroughly by means of shaker can or rotary hand duster. Repeat in 4 to 8 weeks or when necessary. Also treat Foultry Houses as indicated.

**POULTRY HOUSES:** Chicken Red Mite, Northern Fowl Mite, Poultry Lice - Apply thoroughly to roasts, nests, and adjacent areas and to dry floor litter at 1 pound per 50 to 60 sq. ft. of floor surface. Repeat as necessary. Avoid contamination of food and water troughs.

**WARRANTY AND LIMITATION OF DAMAGES:** Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or value, etc.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NET CONTENTS:

EPA Reg. No.: 5905-2

## CAUTION

Read entire label. Use strictly in accordance with cautions and directions, and with applicable state and federal regulations.

Keep out of reach of children. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water. Fish may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Keep pesticide in original containers. Burn bag immediately when empty. Stay out of smoke. Apply thoroughly. Do not apply to food crops within 3 days of harvest unless otherwise specified. This product is formulated for application by ground or air equipment.

### DIRECTIONS FOR USE

**ALFALFA:** Armyworms - 25 lbs. per acre. Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittlebugs, Vetch Bruchids - 30 lbs. per acre. Spotted Alfalfa Aphids - Ground - 12 to 15 lbs. per acre or Air - 12 to 20 lbs. per acre. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the field or are not hanging on outside of hives. May be applied on day of harvest or grazing.

**APPLES:** Red Spider Mites (such as European Red, Two-Spotted, Pacific, Willamette, Clover) (2 or more applications as needed), Green Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphids (1 or more applications as needed) - 30 to 40 lbs. per acre. This product may injure foliage on some varieties of apples.

**APRICOTS:** Spider Mites, Brown Apricot Scale (Crawlers), Orange Tortrix - 50 lbs. per acre. Do not apply within 7 days of harvest.

**ASPARAGUS:** Thrips - 20 to 25 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

**BEANS:** Mexican Bean Beetle, Japanese Beetles, Cucumber Beetles, Leafhopper, Spider Mites, Aphids - 30 to 35 lbs. per acre. Two or more applications as needed. Do not apply within 1 day of harvest.

**BLUEBERRIES:** Japanese Beetles - 20 lbs. per acre. Apply when berries turn blue and at 10-day intervals while infestation persists. Plum Curculio, Sharp-nosed Leafhopper - 32 to 40 lbs. per acre. Do not apply within 1 day of harvest.

**BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, KALE, MUSTARD GREENS, TURNIPS:** Aphids, Imported Cabbageworm, Cabbage Looper, Diamond-back Moth, Flea Beetles - 30 to 50 lbs. per acre. Repeat as necessary. Do not apply to Cauliflower, Brussels Sprouts, Cabbage, Mustard Greens and Kale within 7 days of harvest and to Broccoli and Turnips within 3 days of harvest.

# HELENA

## BRAND

CONS. RELIEF

# 5% CYTHION® DUST

THE PREMIUM GRADE MALATHION®

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL.**

ACCEPTED

10-18-74

2-9065

ACTIVE INGREDIENT:

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC PEACE AND  
ORDER UNDER

By Wt.

Willamette, Clover (2 or more applications as needed), Green Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphids (1 or more applications as needed) - 30 to 40 lbs. per acre. This product may injure foliage on some varieties of apples.

APRICOTS: Spider Mites, Brown Apricot Scale (Crawlers), Orange Tortrix - 50 lbs. per acre. Do not apply within 7 days of harvest.

ASPARAGUS: Thrips - 20 to 25 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

BEANS: Mexican Bean Beetle, Japanese Beetles, Cucumber Beetles, Leafhopper, Spider Mites, Aphids - 30 to 35 lbs. per acre. Two or more applications as needed. Do not apply within 1 day of harvest.

BLEUBERRIES: Japanese Beetles - 20 lbs. per acre. Apply when berries turn blue and at 10-day intervals while infestation persists. Plum Curculio, Sharp-nosed Leafhopper - 32 to 40 lbs. per acre. Do not apply within 1 day of harvest.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, KALE, MUSTARD GREENS, TURNIPS: Aphids, Imported Cabbageworm, Cabbage Looper, Diamond-back Moth, Flea Beetles - 30 to 50 lbs. per acre. Repeat as necessary. Do not apply to Cauliflower, Brussels Sprouts, Cabbage, Mustard Greens and Kale within 7 days of harvest and to Broccoli and Turnips within 3 days of harvest.

CARROTS: Leafhoppers - 30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

COTTON: Boll Weevil, Aphid, Leafworm and Spider Mites - 20 to 30 lbs. per acre. Do not apply within 7 days of hand picking. The time between applications should not exceed 5 to 7 days. If applied by air, wind should be very light. Either morning or evening applications are recommended. Emphasis should be on complete coverage. Consult state agricultural extension service for timing of applications.

CRANBERRIES: Leafhoppers, Black-Headed Fireworms - Ground Machine: 30 lbs. per acre. Make thorough coverage.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites, Pickleworms, Squash Vine Borer - 30 to 35 lbs. per acre. CANTALOUPE - Leafhoppers - 30 to 35 lbs. per acre as necessary. Do not apply unless plants are dry nor within 1 day of harvest.

DATES: Nitidulid Beetles - Apply to cover each cluster thoroughly using 40 lbs. per acre. Make no application on dates within 7 days of the first picking.

GRAPES: Leafhoppers, Spider Mites - 25 to 30 lbs per acre. Leafhoppers - One or more applications as needed. Spider Mites - Two or more applications as needed.

LETTUCE: Aphids, Mites, Cabbage Looper, Leafhoppers - 25 to 30 lbs. per acre. Do not apply to Head Lettuce within 7 days of harvest and to Leaf Lettuce within 14 days.

MOSQUITOES: Apply 10 to 15 lbs. per acre with a power duster. Avoid drift where susceptible fish and wildlife may be endangered.

ONIONS (SEED ONLY): Onion Thrips - 35 to 40 lbs. per acre.

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL.**

ACCEPTED  
10-18-74  
5905-2  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND FUNGICIDE ACT  
FOR ECONOMIC PEST CONTROL  
ED UNDER REG. NO. 100.000  
TO ATTACHED LABEL

ACTIVE INGREDIENT:	Malathion* .....	By Wt.	5%
INERT INGREDIENTS:	.....		95%
	Total .....		100%

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

®Trademark of American Cyanamid Company.

*LOOK UNDER  
HERE*

61174

MANUFACTURED BY  
**HELENA CHEMICAL COMPANY**  
MEMPHIS TENNESSEE

NET CO

**ORNAMENTALS:** Aphids, Red Spider Mites, Whitefly, Thrips, Japanese Beetle, Adult Four-lined Leaf Bug, Bagworm, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper - 25 to 30 lbs. per acre. Repeat as necessary. Do not use on Boston, Maidenhair and Pteris Ferns, and some species of Crassula.

**PEARS:** Pear Psylla, Aphids, Woolly Aphids - 30 to 40 lbs. per acre. One or more applications as needed. Spider Mites - Same directions listed under Apples. Do not apply within 1 day of harvest.

**PEAS:** Pea Aphid - 25 lbs. per acre as needed. Do not apply within 3 days of harvest or 7 days if forage is to be fed to livestock.

**PEPPERMINT, SPEARMINT:** Flea Beetle Adults - 20 lbs. per acre. Do not apply within 7 days of cutting.

**PEPPERS:** Pepper Maggots - 30 lbs. per acre.

**PUMPKINS:** Squash Vine Borers - 35 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply unless plants are dry.

**SPINACH:** Aphids - 25 to 30 lbs. per acre. Do not apply within 7 days of harvest.

**RASPBERRIES:** Aphids, Rose Scale - 35 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

**STRAWBERRIES:** Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Potato Leafhoppers, Strawberry Leafrollers, Whiteflies, Strawberry Root Weevil, Field Crickets - 40 lbs. per acre. Repeat as necessary. Strawberry Root Weevils - Apply to soil surface before planting and work into top 6 to 8 inches of soil.

**TOBACCO:** Aphids - 20 to 30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

**TOMATOES:** Aphids, Spider Mites, Tomato Russet Mite, Drosophila - 35 to 40 lbs. per acre. Do not apply within 1 day of harvest. **MILLIPEDES, SPRINGTAILS, SOWBUGS** In and Around Greenhouses, Around Homes, In Yards and Ornamental Gardens - 1 lb. per 150 sq. ft. (10' x 15') of soil surface or where insects congregate. Repeat at 7 to 10-day intervals as needed.

**FLEAS:** In Dog and Cat Kennels, In Pens, Yards, Lawns, and under Houses - Use 1 to 2 lbs. per 1,000 sq. ft. Use a hand duster. Remove manure or debris before dusting.

**HORNFLIES:** Beef and Dairy Cattle - Apply 3 tablespoonfuls per animal. Do not apply more than 1.5 ounces per Dairy cow. Apply on back and neck of animals. Repeat at 10- to 14- day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not use sprays or dip dairy animals in malathion. Do not contaminate water troughs.

**LICE:** Swine, Beef Cattle - Dust animals thoroughly. Repeat in 10 days and as needed. For pigs less than 1 month of age, use ¼ to ½ tablespoonful per animal. One treatment may be sufficient. Repeat only as needed. Do not treat calves under 1 month old. Avoid contamination of feed, food containers and watering troughs. Do not apply to dairy animals. Swine - Also treat pens thoroughly.

**BEEF AND NONMILKING CATTLE:** Hornflies, Lice, Reduction of Faceflies (Dust-Bag Self Applicators) - Arrange dust-bag self applicators stocked with 10 to 20 lbs. of 5% Cythion Dust at mid-shoulder height near salt boxes or over fenced-in entrances to water troughs, ponds or loafing areas. Mandatory daily use will give the best results. Restock bags as frequently as necessary. Avoid contamination of salt, feed or water. Do not make

**SPRINGTAILS, SOWBUGS** In and Around Greenhouses, Around Homes, In Yards and Ornamental Gardens - 1 lb. per 150 sq. ft. (10' x 15') of soil surface or where insects congregate. Repeat at 7 to 10-day intervals as needed.

**FLEAS:** In Dog and Cat Kennels, In Pens, Yards, Lawns, and under Houses - Use 1 to 2 lbs. per 1,000 sq. ft. Use a hand duster. Remove manure or debris before dusting.

**HORNFLIES:** Beef and Dairy Cattle - Apply 3 tablespoonfuls per animal. Do not apply more than 1.5 ounces per Dairy cow. Apply on back and neck of animals. Repeat at 10- to 14- day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not use sprays or dip dairy animals in malathion. Do not contaminate water troughs.

**LICE:** Swine, Beef Cattle - Dust animals thoroughly. Repeat in 10 days and as needed. For pigs less than 1 month of age, use  $\frac{1}{4}$  to  $\frac{1}{2}$  tablespoonful per animal. One treatment may be sufficient. Repeat only as needed. Do not treat calves under 1 month old. Avoid contamination of feed, food containers and watering troughs. Do not apply to dairy animals. Swine - Also treat pens thoroughly.

**BEEF AND NONMILKING CATTLE:** Hornflies, lice, Reduction of Faceflies (Dust-Bag Self Applicators) - Arrange dust-bag self applicators stocked with 10 to 20 lbs. of 5% Cythion Dust at mid-shoulder height near salt boxes or over fenced-in entrances to water troughs, ponds or loafing areas. Mandatory daily use will give the best results. Restock bags as frequently as necessary. Avoid contamination of salt, feed or water. Do not make accessible to lactating dairy animals or nonlactating animals within 2 weeks of freshening.

**POULTRY:** (Direct Application to Chickens, Geese, Ducks, Turkeys): Northern Fowl Mite, Poultry Lice, Chicken Red Mite (as a supplement to premise treatments for chicken red mite) - Dust individual birds thoroughly by means of shaker can or rotary hand duster. Repeat in 4 to 8 weeks or when necessary. Also, treat Poultry Houses as indicated.

**POULTRY HOUSES:** Chicken Red Mite, Northern Fowl Mite, Poultry Lice - Apply thoroughly to roosts, nests, and adjacent areas and to dry floor litter at 1 pound per 50 to 60 sq. ft. of floor surface. Repeat as necessary. Avoid contamination of food and water troughs.

**WARRANTY AND LIMITATION OF DAMAGES.** Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.