

READ ENTIRE LABEL BEFORE USING



CAUTION

Harmful if swallowed. Avoid breathing of spray mist. Avoid repeated contact with skin. Wash thoroughly with soap and water after using. Keep out of reach of children. Do not use on animals. Do not contaminate feed or foodstuff. This product should not be used for any purpose than that suggested on label

Foris to fishe and wildlife. Keep out of lakes, ponds, and streams. Do nat apply whin tweatier condition favor brieft from areas treated. apply this fraduet only a perification this laber. NOTE TO PHYSICIAN

Cythion®, upon repeated, prolonged emissuse may cause . cholinesterase inhibition Atropine is antidotal. & OAM in alos antidatal

WARRANTY

WARRANTY AND LIMITATION OF DAMAGES. Helene Chemical Company warrants that this methe chemical description on the label and is reasonably fit for the purposes referred te for Use and Conditions of Sale, subject to the inherent risks referred to therein. nical Company makes no other express or implied warranty, including any other express of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company except in writing, with a specific reference to this warranty. Any damages

HELENA

BRAND




5 lb. Emulsion

KEEP OUT OF REACH OF CHILDREN ANNON SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

MANUFACTURED BY

1 gallon of Cyrhion[®] 5 Lb. EMULSION contains 5 lbs. Cythion®



*O.O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate.

SUGGESTIONS FOR USE ON COTTON

HELENA FIVE POUND CYTHION® EMULSION will control the following insects attacking cotton: Boll Weevil, aphids, leafworms, and some species of spider mites.

HELENA FIVE POUND CYTHION® EMULSION contains 5 pounds Cythion® per gallon. Water dilution of various strengths to be used is conventional and low volume sprayers (both ground and airplane applicators) can be prepared. HELENA FIVE POUND CYTHION® EMULSION is mixed with water by agitation or recirculating through spray pumps. If mixture is allowed to stand, re-mix before using-

905

S

b

Cythion[®] sprays are recommended to be at the rate of 1 gal. to 5 acres minimum-I gal. to 3 acres maximum depending upon local conditions. For timing of applications, consult your State Agricultural Extension Service or Experiment Station.

Always use sufficient water for proper coverage.

Do not apply this product where there may be grazing by dairy animals or animals being finished for slaughter.

Do not contaminate feed or foodstuffs. When using this product suitable caution should be followed.

Do not apply this product within seven days of hand picking. Mix with 3-10 gallons water per acre for Aerial Application. Mix with 10-25 gallon water per acre for Ground Application.

COMBUSTIBLE MIXTURE

KEEP AWAY FROM OPEN FLAME


DO NOT USE THIS CONTAINER FOR ANY PURPOSE

U.S.D.A. REG. NO. 5905-7

[®] Trademark Agricultural Division American Cyanamid

HELENA CHEMICAL COMPANY WEST HELENA, ARK. - TAMPA, FLA. - LUBBOCK. TEX.

NET CONTENTS



Roya