

READ ENTIRE LABEL BEFORE USING

WARNING

MAY BE FATAL IF SWALLOWED

Do not breathe spray mist. Do not get in eyes, on skin or clothing. Use may cause cholinesterase inhibition. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. May be fatal if swallowed, inhaled or absorbed through skin.

In case of contact wash immediately with soap and water: for eyes, flush with water for 15 minutes.

Do not contaminate feed or foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash contaminated clothing with soap and hot water before re-use.

STORE OUT OF REACH OF CHILDREN

ATROPINE IS ANTIDOTAL

Commercial operators should keep supply of Atropine 1 - 100 grain tablets.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops.

Toxic to fish and wildlife. Keep out of streams, lakes or ponds. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your State Agricultural Extension Service.

Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes.

The Federal Food and Drug Administration has set official tolerances for certain pesticides. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

DESTROY EMPTY CONTAINER — DO NOT RE-USE

NET CONTENTS

U.S. GALLONS

ACCEPTED
MAY-17-1973
UNDER THE
FUNGICIDE
FOR FOODS
UNDER NO
5905-62

Comments

HELENA

BRAND

TRITHION® 4E INSECTICIDE ACARICIDE

AN ORGANIC PHOSPHATE - EMULSIFIABLE LIQUID

® T.M. Reg. Stauffer Chemical Company for insecticide-acaricide

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL**

1 gallon of
HELENA TRITHION 4-E
contains
4 Lbs. Trithion

ACTIVE INGREDIENTS:

Carbophenothion S (p Chlorophenylthio)	
methyl O,O dimethyl phosphorodithioate	45.2%
Xylene Range Aromatic Hydrocarbon	
Solvent	44.0%

INERT INGREDIENTS:

	10.8%
TOTAL	100.0%

MANUFACTURED BY

HELENA CHEMICAL COMPANY

West Helena, Ark - Tampa, Fla - Lubbock, Texas

Des Moines, Iowa - West Columbia, South Carolina

DIRECTIONS

Pour recommended amounts of the spray tank, add balance of water to the tank during filling, mixing and spray mixture to stand. Do not combine with wettable powders in the same spray tank. Materials being combined has proved to be effective. This product is to be used by

RECOMMENDATIONS

COTTON: Two-spotted mite, carmine mite — Early Season: Use 1/2 - 1 pt. per acre for thorough coverage. Mid and Late Season: Use 1 - 1 1/2 pt. per acre in sufficient water for thorough coverage and repeat applications in 5-7 days. Use 1/2 to 1 pt. per acre in sufficient water for thorough coverage. Cotton Leaf perforator, fleahopper, etc. Use sufficient water for thorough coverage. Do not be grazed or fed to livestock.

SORGHUM: Sorghum midge, two-spotted mite, etc. Use 1 - 1 1/2 pt. per acre in sufficient water for thorough coverage. More than two applications per season within 21 days after application. Use only on some varieties of sorghum. May cause a slight discoloration to the leaves.

WARNING

WARNING AND LIMITATIONS

Helena Chemical Company warrants the chemical description on the label for the purpose referred to in the Director of Pesticide Sale, subject to the inherent risks. Helena Chemical Company makes no other warranty, including any other express or implied warranty of MERCHANTABILITY, and Helena Chemical Company is authorized to do so except in reference to this warranty. Any damage caused by this warranty shall be limited to consequential commercial damage, etc.

EPA Reg No 5905-62

DO NOT RE-USE THIS CONTAINER

ACCEPTED WITH COMMENTS

BEFORE USE

ACCEPTED
MAY-17-1974
UNDER THE FEDERAL
FUNGICIDE ACT
FOR EEDING TO FOLLO
UNDER NO 5905-62

ALLOWED

eyes, on skin or clothing.
For commercial applic-
thing, rubber overshoes and
haled or absorbed through

h soap and water: for eyes.

Wash hands, arms and
before eating or smoking.
and hot water before re-use.

OF CHILDREN

ADULTAL

y of Atropine 1 100 grain

USE PRECAUTIONS

as occupied by unprotected
djoining food and forage

of streams, lakes or ponds.
posed to direct treatment or
ion may be obtained from
ce.

illed. Do not contaminate
posal of wastes.

nisuration has set official
rder that residues on food
lished tolerances, use only
ply closer to harvest than

HEAT OR OPEN FLAME

DO NOT RE-USE

U.S. GALLONS

Comments

HELENA

BRAND

TRITHION® 4E INSECTICIDE ACARICIDE

AN ORGANIC PHOSPHATE - EMULSIFIABLE LIQUID

© T.M. Reg. Stauffer Chemical Company for insecticide-acaricide

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL**

1 gallon of
HELENA TRITHION 4-E
contains
4 Lbs. Trithion

ACTIVE INGREDIENTS:		
Carbophenothion S (p-Chlorophenylthio)		
methyl O O dimethyl phosphorodithioate	45.2%	
Xylene Range Aromatic Hydrocarbon		
Solvent	44.0%	
INERT INGREDIENTS		10.8%
TOTAL		100.0%

MANUFACTURED BY
HELENA CHEMICAL COMPANY
West Helena, Ark Tampa, Fla Lubbock, Texas
Des Moines, Iowa West Columbia, South Carolina

DIRECTIONS FOR USE

Pour recommended amounts of this material into nearly filled spray tank, add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. Do not combine emulsifiable liquids and wettable powders in the same spray tank unless previous use of materials being combined has proved them to be physically compatible. This product is to be used by ground equipment only.

RECOMMENDATIONS FOR COTTON

COTTON: Two-spotted mite, carmine (cinnabar) mite, tropical mite — Early Season: Use 1/2 - 1 pt. per acre in sufficient water for thorough coverage. Mid and Late Sea son: 3/4 - 1 pt. per acre in sufficient water for thorough coverage. If mite populations are high use 1 - 1 1/2 pt. per acre in sufficient water for thorough coverage and repeat applications in 5-10 days. Cotton Leafworm— Use 1/2 to 1 pt. per acre in sufficient water for thorough coverage. Cotton Leaf perforator, fleahopper, aphids—Use 1 qt. per acre in sufficient water for thorough coverage. Treated plants should not be grøzed or fed to livestock.

SORGHUM: Sorghum midge, two-spotted mites—Use 1 pt. per acre in sufficient water for thorough coverage. Do not make more than two applications per season. Do not graze or cut forage within 21 days after application. Under certain climatic conditions and on some varieties of sorghum, organo-phosphorous chemicals may cause a slight discoloration to the leaves.

WARNING

WARNING AND LIMITATIONS OF DAMAGES.
Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

EPA Reg No 5905-62

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE