

**DIRECTIONS FOR USE**  
**VEGETABLES AND SMALL FRUITS**

To control the insects on the crops listed below, apply uniformly and thoroughly to upper and lower surfaces of all foliage. Dust coverage of foliage should be complete. Do not apply in excessive amounts. Begin applications when insects or their damage first appear. Repeat as necessary, generally at 7 day intervals unless shorter interval is stated. May be applied on day of harvest unless otherwise indicated.

**ASPARAGUS** Asparagus beetle. Do not repeat applications within 3 days. Do not apply within 1 day of harvest.

**BEANS** Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetles, Japanese beetle, adult and leafhoppers.

**CORN** Sweet Corn earworm, European corn borer, fall armyworms, flea beetles, Japanese beetle, adults, sap beetle, and leafhoppers. For larvae in whorls, European corn borer, and fall armyworm, and foliage feeders, apply to entire plant before tasseling. For insects attacking silks and ear, apply at 2 to 3 day intervals starting when silks first appear and continuing until they become dry. Three or more applications may be required. Proper timing and coverage are essential.

**CUCUMBERS, MELONS, PUMPKINS AND SQUASH** Pickle worms, melonworms, cucumber beetles, flea beetles, leafhoppers, and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when foliage is wet or during periods of prolonged high humidity. Do not use on watermelons.

**POTATOES, TOMATOES, EGGPLANT, and PEPPERS** Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, tomato fruitworm, tomato hornworm, tarnished plant bug, and stink bug.

**CABBAGE, BROCCOLI, AND CAULIFLOWER** Flea beetles, Harlequin bug, imported cabbage worm, and corn earworm. For cabbage looper, apply at 4 to 5 day intervals. Do not apply within 3 days of harvest. Where cabbage looper is the principal problem, a pesticide that is specifically recommended should be used.

**GARDEN BEETS, LETTUCE, RADISHES, SPINACH, TURNIPS AND CARROTS** Flea beetles, leafhoppers, Harlequin bug, armyworms, Imported cabbage worm, corn earworm, spittlebugs, tarnished plant bug, and stink bugs. For cabbage looper, apply at 4 to 5 day intervals. Do not apply to garden beets, head lettuce, radishes, and turnips within 3 days of harvest. Do not apply to leaf lettuce and spinach or to garden beets and turnips if tops are to be used for food and feed within 14 days of harvest.

**STRAWBERRIES** Meadow spittlebug, strawberry leaf roller, and strawberry weevil. Do not apply within 1 day of harvest.

**GRAPES** Grape berry moth, Japanese beetle, adults, red banded cat flea, leafhopper, and grape leaf skeletonizer.

**BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, and RASPBERRIES** Japanese beetle, adults, leaf rollers, leafhoppers, and raspberry aphid. Do not apply within 7 days of harvest.

**BLEUBERRIES** Blueberry maggot, cherry fruitworm, and cranberry fruitworm. Applications for blueberry maggot should be timed with emergence of adults and repeated according to local schedules.

The rate of application is cubic as uniform and thorough dust coverage of upper and lower surfaces of all foliage is 10 pounds per acre on cucumbers, melons, pumpkins, and squash. 40 pounds per acre on asparagus, beans, French beans, corn, potatoes, tomatoes, eggplant, peppers, cabbage, broccoli, cauliflower, and lettuce, turnips, radishes, spinach, carrots, strawberries, raspberries, blackberries, boysenberries, and raspberries.

# 5% SEVIN<sup>®</sup> DUST

## INGREDIENTS

<b>ACTIVE INGREDIENTS</b>		
Carbaryl (1-naphthyl N-methylcarbamate)	.....	5%
<b>INERT INGREDIENTS</b>	.....	95%
<b>Total</b>		<b>100%</b>

® Trademark Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

See Remainder of CAUTION Statement on this Label.

## CAUTION

This material sold without warranty as to hazards or results to be obtained.

DO NOT REUSE EMPTY CONTAINER. DESTROY WHEN EMPTY.

## CAUTION

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact with the skin. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Do not contaminate feed and foodstuffs.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

MANUFACTURED BY

**HELENA CHEMICAL COMPANY**

WEST HELENA, ARKANSAS

Chemical's Note  
Foot Copy

AGREEMENT WITH COMPANY

w  
l  
f  
o  
th  
w  
o  
r  
is  
w  
l  
d  
a  
N

# SEVIN® DUST

## ORNAMENTAL PLANTS

FLOWERS, SHRUBS, AND SHADE TREES—Apple aphid, bag worm, lace leaf miner, box elder bug, boxwood leaf miner, elm leaf beetle, Japanese beetle, adults, lace bugs, leafhoppers, leaf roller, mealy bug, mimosa webworm, periodical cicada, rose aphid, rose slug, tent caterpillar, and thrips (exposed) on ornamental plants and shade trees such as rose, carnation, chrysanthemum, gladiolus, lilac, arbor vitae, juniper, hydrangea, azaleas, elm, maple, oak, dogwood, birch, and pine. Apply thoroughly and uniformly to upper and lower surfaces of all foliage. Repeat application at 7 day intervals as needed. Do not apply to Boston ivy.

## POULTRY

CHICKENS, PARTRIDGES, PHEASANTS, AND PIGEONS—Northern fowl mite, chicken red mite, lice, fleas, and bed bugs in houses or quarters. Apply 1 pound per 40 sq. ft. of floor and litter, and to roosts and interior surfaces. Do not apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of slaughter.

For Northern fowl mite and lice on the birds, as a supplement to treatment of premises, apply 1 pound per 100 birds. Direct dust to vent and fluff areas, and distribute throughout feathers. Do not repeat within 4 weeks. Do not apply within 7 days of slaughter.

DO NOT CONTAMINATE FEED, DRINKING WATER, FEEDING AND WATERING RECEPTACLES.

## PET ANIMALS

DOGS AND CATS—Ticks and fleas. Apply liberally over entire animal, avoiding eyes, and rub thoroughly into hair. Comb out dead ticks several hours after treatment. Repeat at weekly intervals as needed. Do not apply to kittens or puppies under 4 weeks of age.

For fleas and ticks in quarters used by dogs and cats, apply liberally into and around sleeping quarters, kennels, or houses. Force dust into cracks and crevices of floor and walls. Repeat as necessary.

DO NOT CONTAMINATE FEED OR FEEDING AND WATERING RECEPTACLES.

## SPECIAL NOTES

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application; does not control spider mites, but is compatible with most commonly used miticides.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

E.P.A. REG. NO. 5905-48

**WARRANTY AND LIMITATIONS OF DAMAGES.** Helena Chemical Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

NET WEIGHT

5.905 95  
**ACCEPTED**

POUNDS

## INGREDIENTS

INGREDIENTS	5%
(1-naphthyl N-methylcarbamate)	95%
INGREDIENTS	100%

Union Carbide Corp. for 1-naphthyl N-methylcarbamate.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

Read remainder of CAUTION Statement on this Label.

## CAUTION

Material sold without warranty as to hazards or results to be expected.

USE EMPTY CONTAINER DESTROY WHEN EMPTY

## CAUTION

If inhaled or swallowed. Do not breathe dust. Do not take food or drink. Contact with eyes or skin may be harmful. Avoid contact with the skin. Wash hands before eating. Take shower or bath after work. Wear regular work clothing. Change to clean clothing daily. Do not contaminate foodstuffs.

Carbamyl is a moderate, reversible, cholinesterase inhibitor. It is antidotal.

MANUFACTURED BY

HELENA CHEMICAL COMPANY

WEST HELENA, ARKANSAS