

**CAUTION:** Handle caustic soda (household lye) with extreme care. Do not get on skin, in eyes or on clothing. Read and carefully follow the precautions on package. Punch holes in top and bottom of container. Crush container. Bury deeply in an isolated area.

**IF THE ABOVE RINSE METHOD CANNOT BE USED**

Punch holes in TOP of container and burn in hot fire until all paint has been completely burned off container. Extreme care should be exercised to insure that containers are completely empty and that persons stay well away from smoke and fumes. Do not attempt to burn more than 5 containers at one time.



**ALFALFA:** Flea beetles, leafhoppers, climbing cutworms, armyworms - Apply 1/2 - 1 pint per acre. Aphids (including pea aphid and spotted alfalfa aphid) - Apply 1/2 - 1 pint per acre. For spotted alfalfa aphid, apply early in spring. Alfalfa weevil (larvae) - Apply 1/2 - 1 pint per acre. Apply when 75% terminals show feeding, when buds and shoots appear ragged, or when the larvae count reaches 25 per sweep. Lygus bugs, mites, stink bugs - Apply 1/2 - 1 1/2 pints per acre. Webworms, clover leaf weevil, alfalfa caterpillar, alfalfa weevil (adult) (Refer also to Egyptian weevil) thrips, and grasshoppers - Apply 1 pint per acre. Egyptian alfalfa weevil - Apply 1-2 pints per acre. Do not apply within 15 days of harvest. Alfalfa seed Chalcid - Apply 1 1/2 - 2 1/2 pints per acre. Do not apply within 20 days of harvest.

**ALMONDS:** Peach twig borer, San Jose scale, parlatoria scale, mite eggs - Apply 2 - 4 pints per acre. For dormant application use 2-4 pints plus dormant oil emulsion or dormant emulsive oil at the dosage recommended for the specific oil product per 100 gals. Fruit tree leafroller, tent caterpillar, peach twig borer - Apply 2 - 4 pints per acre (per 100 gals. of water). Do not apply after hulls open. Do not exceed 600 gals. dilute spray per acre.

**APPLE:** Aphids (including apple, rosy apple and woolly apple aphid), European red mites - Apply 1/2 - 1 pint per acre (per 100 gals. water). Scales, redbanded leafroller - Apply 1 - 2 pints per acre (per 100 gals. of water). Codling moth, plum curculio - Apply 1 - 2 pints per acre (per 100 gals. water). Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 15 days of harvest. Fruit and foliar injury may occur on McIntosh, Cortland, and related varieties. Consult your State Cooperative Agricultural Extension Service for further information.

**APRICOTS:** Aphids (including green peach aphid), European red mites - Apply 1/2 pint per acre (per 100 gallons of water). Do not apply within 14 days of harvest.

**ARTICHOKES:** Aphids, armyworms (to 3rd instar), plant bugs - Apply 1 pint per acre. Cutworms (including climbing and surface feeding species), cabbage looper, artichoke plum moth - Apply 2 pints per acre. If plants are cut in April and May, begin applications after September and repeat at 14 to 21 day intervals as needed through December. 30 day intervals through January and February, 14 day intervals from March through the end of picking season. Workers should not re-enter fields for 7 days following treatment. Do not apply within 7 days of harvest.

**BARLEY:** Thrips - Apply 1/2 - 1/2 pints per acre. Mites (brown wheat), leafhopper, cutworms (climbing) - Apply 1/2 to 1 pint per acre. Stink bugs, armyworm (to 3rd instar), black grass bugs, aphid (including greenbug) - Apply 1/2 - 1 1/2 pints per acre. Apply in early spring for green bug control. Thrips, Say's stink bugs, spider mites, false chinch bug, grasshopper, chinch bugs - Apply 1 1/2 pints per acre. Do not apply within 15 days of harvest.

**BEANS:** Leafhoppers, flea beetles, plant bugs (including Lygus bugs), aphids, armyworms - Apply 1 - 2 1/2 pints per acre. Cabbage looper - Apply 2 pints per acre. Stink bugs, cutworms (including climbing and surface feeding species) - Apply 2 - 2 1/2 pints per acre. Do not apply within 15 days of harvest. For green and lima beans do not apply within 21 days of harvest for dosage rates exceeding 1 pint per acre.

**BEETS:** Mites, aphid, armyworms, flea beetles, leafhopper - Apply 1 - 2 pints per acre. Stink bugs, plant bugs (including Lygus bugs), cabbage looper - Apply 2 pints per acre. Do not apply within 15 days of harvest. Do not apply within 21 days if tops are to be used for food or feed.

**BROCCOLI:** Thrips - Apply 1/2 - 1 pint per acre. Aphids, armyworms - Apply 1/2 - 2 pints per acre. Flea beetles, leafhopper, mites - Apply 1 - 2 pints per acre. Imported cabbageworms, cabbage looper, stink bugs - Apply 2 - 2 1/2 pints per acre. Plant bugs - Apply 2 1/2 pints per acre. Do not apply within 21 days of harvest.

**BRUSSEL SPROUTS:** Aphids, armyworms, thrips - Apply 1/2 - 2 pints per acre. Flea beetles, leafhoppers, mites - Apply 1 - 2 pints per acre. Imported cabbageworms, cabbage looper, stink bugs - Apply 2 - 2 1/2 pints per acre. Plant bugs - Apply 2 1/2 pints per acre. Do not apply within 1 days of harvest.

**CABBAGE:** Aphids, armyworms, thrips - Apply 1/2 - 2 pints per acre. Flea beetles, leafhopper, mites - Apply 1 - 2 pints per acre. Imported cabbageworms, cabbage looper, stink bugs - Apply 2 - 2 1/2 pints per acre. Plant bugs - Apply 2 1/2 pints per acre. Do not apply within 21 days of harvest.

BEST AVAILABLE COPY

90812

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

STOP - READ LABEL BEFORE USING

**POISON**

See Side Panels  
for Antidote &  
Precautions




**DANGER**

Keep Out of  
Reach of  
Children



**4 LB. Methyl Parathion**  
Emulsifiable Insecticide Concentrate



1 gallon of  
**HELENA 4 LB**  
**METHYL PARATHION**  
contains 4 pounds of  
Methyl Parathion

**ACTIVE INGREDIENTS (by weight)**  
\*O,O-dimethyl O-p-nitrophenyl  
phosphorothioate 45.6%  
Xylene 47.4%  
**INERT INGREDIENTS (by weight)**  
TOTAL 7.0%  
100.0%

\*Methyl Parathion

EPA REG. NO. 5905-55

EPA REG. NO. 5905-55

**NOT FOR HOME USE**