

WARNING

1. This product is toxic and is absorbed through the skin in dry form and in solutions. Do not breathe spray mist. Do not get in eyes or on skin. Use U.S. Bureau of Mines approved respirator or goggles if necessary. May be fatal if swallowed.
2. Keep out of reach of children.
ANTIDOTE: Internal: Give one tablespoonful of 1% solution of granules to drink mixed in glass of warm water. Repeat until vomit is clear. Call doctor immediately.
External: Wash thoroughly with soap and water.
3. In case of spillage on skin, wash with soap and water.
4. Avoid contamination of feed and foodstuffs.
5. Keep clothing free from spray residue.
6. Do not spray on pets.
7. Do not apply to plants belonging to the Cucurbitaceae (cucumber, squash, melon, etc.).
8. Do not feed plants which have been treated with this product to dairy animals or to animals being finished for slaughter.
9. Avoid spraying plants in bloom when bees are active in the field. Do not spray or make applications late in the evening when bees are foraging.
10. Do not use this product in lawns. Birds feeding on treated grass may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

HELENA

6 LB. Toxaphene


Emulsifiable Insecticide Concentrat

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE P/

1 gallon of
HELENA 6 LB
TOXAPHENE
contains the equivalent of
6 pounds Toxaphene
per gallon

ACTIVE INGREDIENTS (by weight)
• Toxaphene
Xylene
INERT INGREDIENTS (by weight)
TOTAL



ANTIDOTE: Internal: Give one tablespoonful of 1% solution of sodium sulfide in a glass of warm water. Repeat until vomit is clear. If ill again, repeat immediately.
External: Wash thoroughly with soap and water.

3. If on clothing, remove clothing, wash with soap and water.
4. Avoid contact with eyes. If contact occurs, wash with water.
5. Keep children away from spraying area.
6. Do not spray around pets.
7. Do not apply to plants belonging to those who do not use (cucumber, squash, melon, etc.).
8. Do not feed plants which have been treated with this product to dairy animals or to animals being finished for slaughter.
9. Avoid spraying plants in bloom when bees are active in the field. If necessary to spray, make applications late in the evening when bees are less active.
10. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NON WARRANTY NOTICE

6 LB. Toxaphene


Emulsifiable Insecticide Concentrat

WARNING

**KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE P/**

1 gallon of
HELENA 6 LB
TOXAPHENE
contains the equivalent of
6 pounds Toxaphene
per gallon

ACTIVE INGREDIENTS (by wei
• Toxaphene
Xylene
INERT INGREDIENTS (by weight
TOTAL




LENA

Toxaphene Insecticide Concentrate

KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS ON SIDE PANEL

ACTIVE INGREDIENTS (by weight)	
Toxaphene	60.0%
Xylene	12.0%
INERT INGREDIENTS (by weight)	8.0%
TOTAL	100.0%



SUGGESTIONS FOR USE

ON COTTON

HELLENA 6-Lb. Toxaphene Emulsion is suggested for following insects on cotton: Bollworm, Fleahopper, Thrips, Grasshopper, Root Borer, Turned Root Borer and moderate infestation of bollworm.

HELLENA 6-Lb. Toxaphene Emulsion is suggested for control of bollworm on cotton. Aerial application of emulsion is suggested for control of bollworm on cotton. Aerial application of emulsion is suggested for control of bollworm on cotton. Aerial application of emulsion is suggested for control of bollworm on cotton.

HELLENA 6-Lb. Toxaphene Emulsion is suggested for control of bollworm on cotton. Aerial application of emulsion is suggested for control of bollworm on cotton. Aerial application of emulsion is suggested for control of bollworm on cotton.

Apply 2 to 3 pounds actual toxaphene per acre, use 2 2/3 pints to 4 pints of product diluted in sufficient water to give coverage. For timing of applications, consult your local representative of extension service or experiment station and repeat applications as local conditions warrant.

For aerial application use with 1-3 gals. of water per acre.


Do not apply this product where there may be grazing livestock or where animals are being fed. Do not apply to any area where livestock are being fed. Do not apply to any area where livestock are being fed.

Do not apply this product to any area where livestock are being fed. Do not apply to any area where livestock are being fed.

DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE

COMBUSTIBLE MIXTURE

Technical Chlorinated Camphene (Chlorine
content 40-50%)



PRODUCED BY
CAL COMPANY
TENNESSEE

This product toxic to fish and wildlife. Do not apply to any area not specified on the label.

DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

COMBUSTIBLE MIXTURE

KEEP AWAY FROM OPEN FLAME

E.P.A. Reg. No. 5905-56

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE