DIRECTIONS

COTTON Defoliation Apply [illegible] date. Do not treat later than [illegible] of the plant [illegible] [illegible] MACHINE [illegible] make [illegible] are [illegible] water [illegible] of the [illegible] soil moisture [illegible] may [illegible] factory [illegible] lack of [illegible] graze treated [illegible] feed [illegible]

GRAIN SORGHUM: As a desiccant apply 1 1/3 to [illegible] gallons per acre in 12 to 15 gallons of water 7 to 10 days before harvest.

MIXING: Agitate Helena 3 LB. Chlorate Defoliant - Desiccant in water until completely dissolved. Use mechanical agitator.

NOTICE

Do not allow spray to drift [illegible] other plant growth that might be defoliated. Clean and wash spray equipment before and after use and do not allow wash water to contact desirable plants. Do not store near foodstuff.

CAUTION: Harmful if swallowed. Avoid inhalation. Avoid skin contact.

CAUTION

Do not allow man or animal to take internally. Wash clothing after handling. Do not contaminate feed and foodstuffs. Do not apply in any manner not specified on label. Do not graze treated areas or feed gin waste to livestock.

Do not contaminate water by cleaning of equipment or disposal of wastes.

HELENA T.M.

3 LB. CHLORATE DEFOLIANT - DESICCANT

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

1 gallon of
HELENA 3 LB.
CHLORATE DEFOLIANT - DESICCANT
contains 3 pounds of
Sodium Chlorate
with a fire retardant

ACTIVE INGREDIENT:	
Sodium Chlorate	27.50%
INERT INGREDIENTS	72.50%
TOTAL	100.00%

GENERAL

Weather conditions and plant cor
ditions that favor good defoliation a
wind velocity, high to adequate so
good defoliation are ample fertility
Nutrients should be ample but not t
GREEN AND TURGID (NOT WIL
LACK OF FERTILIZERS).

MATURITY - Boll load heavy, all
little or no second growth. Condit
are those adverse to the above.

ACCEPTED
March 7, 1975
UNDER [illegible] FOR ECONOMIC [illegible] REGISTERED UNDER NO. 5905-118 SUBJECT TO ATTACHED COMMENTS.

WA

[illegible]

EPA [illegible] 5905-118