

 **POISON** 

**DANGER! POISONOUS BY SKIN CONTACT
INHALATION OR SWALLOWING.**

RAPIDLY ABSORBED THROUGH SKIN

**REPEATED EXPOSURE MAY, WITHOUT
SYMPTOMS, BE INCREASINGLY HAZARDOUS**

Do not get in eyes, on skin, on clothing.

Do not breathe spray mist or vapors.

Wear clean, heavy rubber gloves and clean or freshly
laundered protective clothing and goggles when applying
or handling. Destroy and replace gloves frequently.

Wear an effective mask or respirator of a type passed by
the U. S. Bureau of Mines for EPN protection.

Do not contaminate feed or foodstuffs.

Wash hands, arms, and face thoroughly with soap and
warm water before eating or smoking.

Bathe immediately after work and change all clothing.

Wash clothing thoroughly with soap and hot water be-
fore re-use.

In case of contact, immediately remove contaminated
clothing and wash skin thoroughly with soap and hot
water; for eyes, flush with water for 15 minutes and get
medical attention.

ANTIDOTES


INTERNAL: Give a tablespoonful of salt in a glass of
warm water and repeat until vomit fluid is clear.

EXTERNAL: Immediately remove all contaminated
clothing and wash thoroughly with soap and warm



2 LB. EPN

EMULSIFIABLE INSECTICIDE CONCENTRATE

DANGER  **POISON** **KEEP OUT OF REACH OF CHILDREN**
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

1 gallon of
HELENA
2 Lb. EPN
contains
2 Lbs. EPN

ACTIVE INGREDIENTS:
O-Ethyl O-p-nitrophenyl Phenyl
phosphonothioate 24.914%
Aromatic Petroleum Solvent 71.013%
INERT INGREDIENTS 4.073%
TOTAL 100.000%
*EPN - A DuPont Product

ACCEPTED

AUG 9 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 5705-10/SUBJECT

DOSAGE AND PE
Apply to fruiting co
control of:

Cotton Fleahopper
Yellow Striped Arm
Spider M

BOLLWEEVIL CO
WHEN TO APPLY

Make weekly inspect
apply HELENA

Boll Weevil: Begin
ing freely and 10 to

Repeat at 5-day interv

OTHER PESTS AS
lications when species
numbers (consult Sta

Repeat at 5-day interv

Aircraft Application—

Ground Application—

This product is toxic
Do not contaminate
any area not specified

Do not apply when we
areas treated. Do not
equipment or disposal

IN
Keep animals and unp
ment and while there

DOSAGE AND PESTS CONTROLLED:
Apply to fruiting cotton at rate of 1.25 pts. per acre for control of:

Cotton Fleahopper	Cotton leafworm
Yellow Striped Armyworm	Thrips
Spider Mites (Certain Species)	

BOLLWEEVIL CONTROL: apply at 2 pts. per acre.

WHEN TO APPLY:

Make weekly inspections to observe infestation, and apply HELENA 2 lb. EPN as indicated below.

Boll Weevil: Begin applications when cotton is fruiting freely and 10 to 25% of squares are punctured.

Repeat at 5-day intervals as long as needed.

OTHER PESTS AS LISTED ABOVE : Begin applications when species appear in potentially damaging numbers (consult State Agricultural Extension Service).

Repeat at 5-day intervals as long as needed.

Aircraft Application—1-3 gal. water per acre.

Ground Application—2-5 gal. per acre.

This product is toxic to fish, birds, and other wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

IMPORTANT

Keep animals and unprotected persons away during treatment and while there is danger from drift or residues. Do not apply within 3 days of picking. Do not permit grazing of areas which have received late applications. Destroy empty containers by burning or burying. Stay out of smoke and vapors.

EPA REG. NO. 5905-101

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE