

## FIRST AID

ATROPINE IS AN ANTIDOTE — CONSULT PHYSICIAN FOR EMERGENCY SUPPLY. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

## ANTIDOTES

**INTERNAL**—Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

**EXTERNAL**—Immediately remove all contaminated clothing and wash skin thoroughly with soap and warm water.

**NOTE TO ATTENDING PHYSICIAN**—Repeat therapeutic doses of atropine may be effective. Do not use morphine.

**STORE IN A COOL, DRY PLACE, AWAY FROM FEED, FOODSTUFFS AND DOMESTIC ANIMALS. KEEP AWAY FROM HEAT AND OPEN FLAME. KEEP CONTAINER CLOSED.**

Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water; for eyes, flush with water for 15 minutes and get medical attention.

## SYMPTOMS OF POISONING

The symptoms of poisoning by organophosphorus insecticides including EPN, are likely to be as follows:

- 1—Digestive upsets, nausea, vomiting, cramps, feeling of pressure under breast bone, diarrhea.
- 2—Sweating and excessive salivation.
- 3—Restlessness, giddiness, headache.
- 4—Muscle tremors, twitching of the facial and eye muscles.

If any of these symptoms are felt, discontinue operations at once; remove person from area; give two atropine tablets (each 1/100 gr.) and CALL A PHYSICIAN. Do not resume with ANY organophosphorus insecticide, including EPN, until the physician

# HELENA BRAND

## 2% EPN INSECTICIDE GRANULES

For Control Of  
European Corn Borer



KEEP OUT OF REACH OF CHILDREN  
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

See Additional Cautionary Information on Side Panel

### ACTIVE INGREDIENT

O-Ethyl O-p-nitrophenyl

phenylphosphonothioate ..... 2%

INERT INGREDIENTS ..... 98%

Total ..... 100%

## DIRECTIONS

HELENA BRAND 2% EPN Insecticide is recommended for use on corn to control European corn borer. Apply at rate of 25 lbs. per acre, using standard granular insecticide application equipment.

**SWEET CORN**—Make first application when 25% of the plants show evidence of leaf feeding and continue at 5-day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch.

**FIELD CORN**—Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present. For control of second brood borers, apply when eggs begin to hatch. For control of first brood cornborers, apply 2% EPN 10 days after the eggs first hatch or when 75% of the plants show feeding damage in the leaf whorls.

## IMPORTANT

Do not apply within 14 days of harvest. Keep animals and unprotected persons away during treatment and while there is danger from drift or residues.

This product will kill fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. (Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.)

Do not reuse this container. Crush and bury when empty. Bury spillage; clean up area with strong lye solution.

**DANGER!** Extremely Hazardous By Skin Contact, Inhalation, or Swallowing. Rapidly absorbed through Skin. Exposure may, without symptoms, be increasingly hazardous.

Do not get in eyes, on skin, on clothing. Do not breathe vapor or spray mist, wear clean heavy, natural rubber gloves, rubber boots, and clean waterproof or freshly laundered protective clothing (coveralls, caps), and goggles when applying or handling. Destroy and replace gloves frequently. Wear an effective mask or respirator of a type passed by the U. S. Bureau of Mines for EPN protection. Do not contaminate feed or foodstuffs.

**WARRANTY AND LIMITATIONS OF DAMAGES.** Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions for Sale, subject to inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including