



RED →

RED ↓

MAY BE FATAL IF SWALLOWED OR ABSORBED THROUGH THE SKIN

MARAUDER

WEEDS IN SOYBEANS AND PEANUTS

ate 21.2
-butyl-4
10.9
67.9
100.0

lphtr

l

nds of a
ate an

KEEP FROM FREEZING
SHAKE WELL BEFORE USING
STORE IN A COOL DRY LOCATION
DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

HELENA BRAND MARAUDER is a selective herbicide combination for the control of the following annual grasses and broadleaf weeds:

ANNUAL GRASSES: foxtail, crabgrass, Barnyardgrass, goosegrass, stinkgrass, sandbur, watergrass, sponge grass, windmill grass, cupgrass, and Johnson grass (from seed).

BROADLEAF WEEDS: Florida pursley, charweed, lamb quarters, ragweed, purslane, velvetleaf, one year bindweed, ground cherry, mustard, shepherd's purse, pigweed, quackweed, arpetweed, morning glory, beardweed, thistle, and teaweed (except yellow).

PREPARATION OF SOIL: The area to be treated should be well prepared and as free as possible from old weeds. If the land has been plowed, the soil should be broken up by harrow or roller wheel. Planting should be promptly resumed. If weather factors make replanting necessary, after **HELENA BRAND MARAUDER** has been applied, thoroughly disk the area prior to replanting.

METHOD OF APPLICATION: Thorough and even coverage of the soil surface is essential for good control. Apply **HELENA BRAND MARAUDER** as a spray at planting time using either a planter mounted sprayer or a ground sprayer after the planting operation. The herbicide should be applied up to the time of crop emergence or on peanuts put before the crop plants emerge.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this product conforms to the chemical description on the label and is substantially fit for the purposes referred to in the Directions and Conditions of Sale, subject to the inherent limitations thereon.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except with specific reference to this warranty. Any damages from a breach of this warranty shall be limited to direct damages and shall not include consequential or other damages, loss of profit or value of crop.

MPANY
7

EPA Reg. 5905-173

DAER

PEANUTS

METHOD OF APPLICATION: For uniform and firm coverage of the soil surface is essential for weed control. Apply **HELENA BRAND MARAUDER** at a steady planting time using either a planter mounted sprayer or a separate sprayer after the planting operation. The chemical may be applied at the time of crop emergence or at any time thereafter but before the crop plants emerge in soybeans.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes neither express nor implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANDISABILITY**, and no agent of Helena Chemical Company is authorized to do so, except in writing, with specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and not included consequential commercial damages which may be incurred.

EPA Reg. 5905-173



DO NOT GET IN EYES, ON SKIN, ON CLOTHING. DO NOT BREATHE SPRAY DRIFT. DO NOT TAKE INTERNALLY. DO NOT WEAR CONTAMINATED CLOTHING OR SHOES. KEEP AWAY FROM HEAT AND OPEN FLAME. NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

ANTIDOTE AND FIRST AID TREATMENT

SYMPTOMS OF POISONING: Irritation, fatigue, sweating, thirst and fever. If symptoms develop, develop from any type of exposure. **SEND FOR A PHYSICIAN.**

FIRST AID: Have patient lie down and rest. Give ample liquid. If feverish, give plenty of liquid. If symptoms persist, consult a physician.

ANTIDOTE: If swallowed, **SEND FOR A PHYSICIAN.** Induce vomiting by giving an emetic. Give a tablespoonfuls of table salt in a glass of warm water, repeat until vomit has cleared. Then give two brandies of brandy in a glass of water. Repeat. **FIRST AID.**

IF SPLASHED IN EYES: Immediately flush eyes with plenty of water. Get medical attention as soon as possible.

IF SPILLED ON SKIN: Immediately wash with plenty of water. **INCLUDING SHOES.** Wash with plenty of water. **FIRST AID.**

NOTE TO ATTENDING PHYSICIAN: A toxicologist should be consulted. Amount of chemical ingested.

DIRECTIONS FOR USE: For use on peanuts, recommended amount of water in the spray tank and the required amount of **HELENA BRAND MARAUDER** must be determined by the water in the tank at the time of application. For use on soybeans, see the label on the container.

PEANUTS

HELENA BRAND MARAUDER is a selective herbicide for the control of weeds in peanuts. It is applied in a spray tank. The amount of water in the tank at the time of application is important. For use on peanuts, see the label on the container.

RECOMMENDED DOSAGE RATE FOR PEANUTS
10 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre
12 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre
14 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre

AMOUNTS OF MARAUDER PER ACRE TO USE FOR BAND TREATMENT OF PEANUTS

WIDTH OF BAND	ROW SPACING
12	40" 50" 40" 40" 40" 40"
14	40" 50" 50" 50" 50" 50"

Use 1/2 lb. per acre of water per acre. Plant in a band 1/2 row wide. Water in the band. Apply the chemical in the band. Water in the band. Apply the chemical in the band.

SOYBEANS

HELENA BRAND MARAUDER is a selective herbicide for the control of weeds in soybeans. It is applied in a spray tank. The amount of water in the tank at the time of application is important. For use on soybeans, see the label on the container.

RECOMMENDED DOSAGE RATE FOR SOYBEANS

10 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre
12 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre
14 - **HELENA BRAND MARAUDER** 4.0 lbs. per acre

AMOUNT OF MARAUDER PER ACRE TO USE FOR BAND TREATMENT ON SOYBEANS

WIDTH OF BAND	ROW SPACING
12	40" 50" 40" 40" 40" 40"
14	40" 50" 50" 50" 50" 50"

Use 1/2 lb. per acre of water per acre. Plant in a band 1/2 row wide. Water in the band. Apply the chemical in the band. Water in the band. Apply the chemical in the band.

HELENA BRAND MARAUDER is a selective herbicide for the control of weeds in soybeans. It is applied in a spray tank. The amount of water in the tank at the time of application is important. For use on soybeans, see the label on the container.

DO NOT USE MARAUDER ON SILT LOAM SOILS OF EXTREMELY FINE TEXTURE

CONTAINER STORAGE AND HANDLING: **HELENA BRAND MARAUDER** is a highly volatile liquid. It may be flammable in the presence of heat, sparks, or flames. It is a strong irritant to the eyes, nose, and throat. It is a strong irritant to the skin. It is a strong irritant to the respiratory system. It is a strong irritant to the mucous membranes.

CONTAINER DISPOSAL

Do not reuse this container for any other purpose. Do not dump this container into the ground. Do not burn this container. Do not incinerate this container.

EQUIPMENT CLEANING

Do not use this container for any other purpose. Do not dump this container into the ground. Do not burn this container. Do not incinerate this container.

DANGER!

Do not reuse this container for any other purpose.