HELENA CHEMICAL COMPANY

1076



DEST have been been

CORPORATE OFFICE

Suite 3200 — Clark Tower 5100 Poplar Avenue Memphis, Tennessee 38137 Telephone: 901/761-0050 Telex. 8105911595

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification

HELENA

EPN 4-E


EMULSIFIABLE INSECTICIDE CONCENTRATE

ACTIVE INGREDIENTS

*O-Ethyl O-(p-nitrophenyl) phenylphosphonothioate	45.87%
Xylene	43.56%
INERT INGREDIENTS.	10.57%
TOTAL	100.00%

*EPN

This product contains 4 lbs EPN per gallon



PRECAUCION AL USARIO: Si usted in lee ingles, no use este producto hasta que le etiqueta haja sido explicado ampliamente. (TRANSLATION: TO THE USER: If you cannot read English, do att use this product until the label has been fully explained to you.)

EPA Reg. No. 5905-174 EPA Est. No. * Net Contents

. . ' . .

Manufactured by Helena Chemical Company Momphis, TN 38137 ACCEPTED

NOV 0 4 1983 Under the Fadaral Insecticida. innaicide icida Act ه ارز **የ**ፅ ና

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - POISON

Poisonous if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Repeated inhalation or skin contact may, without symptoms progressively increase susceptibility to EPN poisoning.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

LYE CONTACT - Flush immediately with clean water for 15 minutes. Get medical attention.

IF BREATHED - Remove victim to fresh air. Get medical attention.

SKIN CONTACT - Remove contaminated clothing and immediately wash skin with soap and water. Get medical attention.

POISONING SYMPTOMS: Weakness, headache, blurred vision, nausea, vomiting, diarrhea, cramps, non-reactive pinpoint pupils, salivation, sweating and discomfort in chest.

NOTE TO PHYSICIAN (Treatment): Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to 'ry small doses of any cholinesterase inhibitor. Allow no further exposure until the time for cholinesterase regeneration has been determined by blood tests. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drive from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blocming crops or weeds. Do not apply this product or allow it to drift to blocming crops or weeds if bees are visiting the treatment area.

PHYSTCAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

9M12 30FC

RE-ENTRY STATEMENT AND WORK SAFETY RULES

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Do not enter treated areas without protective clothing until sprays have dried.

I cause certain states may require more restrictive reentry intervals for various crops treat ϵ 1 with this product, consult your State Department of Agriculture for further information.

PROTECTIVE CLOTHING REQUIRED: Wear clean protective clothing, goggles, and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within 24 hours of treatment.

The following protective clothing must be worn: lightweight, unlined, natural rubber gloves at least midforearm in length; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a nonpermeable, non-cloth material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least midcalf in length; full-face respirators and goggles are required. Aerial applicators in positive pressure cockpits and other applicators in comparable ground equipment with appropriate filters at all air intakes need not comply with these protective clothing requirements.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warr which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: '(Appropriate signal word DANGER or WARNING). Area treated with (name of pesticide) on (date of application). Do not enter without appropriate protective clothing for 24 hours. (Insert here actions to take in case of accidental exposure.)'

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide disposal: Pesticide wastes are acutely hazardous. Improper disposal of ":": excess resticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container disposal: Triple rinse (or equivalent). Then offer for recycling or recording ioning, or puncture and dispose of in a sanitary landfill, or by other procedures ... approved by state and local authorities.

- - our 1

USE DIRECTIONS

Do not apply this product when weather conditions favor drift from treated area. When applying this product use sufficient spray volume to thoroughly cover all surfaces to be treated. Water emulsions may be easily applied with either ground equipment or by airplane for the control of many insects. For AERIAL APPLICATION, mix with 1 to 3 gallons of water. (For California use a minimum of 5 gallons.) For aerial applications, flagging must be by fully automated mechanical means or by humans working in totally enclosed vehicles. For GROUND APPLICATION, mix with approximately 3 to 10 gallons of water per acre. (For California use 5 to 20 gallons.)

Do not use on cucurbit crops (Cucumbers, Cantalopes, Melons, Squash) as injury to plants may result.

(OPTION: Do not apply to blooming cotton if bees are visiting the treatment area.

FOR CONTROL OF COTTON FLEAHOPPER, COTTON LEAFWORM SPIDER MITES THRIPS, YELLOW STRIPED ARMYWORM — 5/8 to 7/8 pint per acre Repeat at 4 to 5-day intervals

BOLLWEEVIL --- 1 to 2 pints. Apply when plants are fruiting freely and 10 to 25% of the squares are punctured. Repeat at 5-day intervals.


BOLLWORM, TOBACCO BUDWORM – 2 pints for Bollworm. Apply when eggs or 4 to 5 larvae per 100 terminals are found. Repeat at 5-day intervals.

Do not apply within 3 days of hand picking. Workers entering fields within 24 hours of application should wear protective clothing. Do not feed treated forage or gin trash to livestock.

BEANS: For control of TWC SPOTTED SPIDER MITE and MEXICAN BEAN BEETLE -- Apply 1 pint per acre. Do not apply will hin 21 days of harvest. (For Mexican Bean Beetle make 2 to 4 applications starting 5 to 7 days after eggs are laid.)

CORN: For control of EUROPEAN CORN BORER – A \rightarrow at rate of 1 pint per acre. On Sweet Corn make application when 25% of the plants show evice, ce of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From 1 to 3 treatments may be required. On Field Corn apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding leaf whorls. For second brood apply when eggs begin to hatch. For corn grown for seed, apply when 25% of plants have feeding damage in the leaf whorls. Repeat in 7 days if new feeding — image occurs. For second brood, apply all pollen shedding and while plants are in the green site... age. Do not apply within 14 days of har is st

Do not apply to corn during the pollen shed periof if bees are visiting the treatment area.



- 4र्ज ।

:.

....

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

THE DIRECTIONS FOR USE OF THIS PRODUCT REFLECT THE OPINION OF EXPERTS BASED ON FIELD USE AND TESTS. THE DIRECTIONS ARE BELIEVED TO BE RELIABLE AND SHOULD BE FOLLOWED CAREFULLY. HOWEVER, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENILY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS OR OTHER UNINTENDED CON: QUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF HELENA OR THE SELLER. ALL SUCH RISKS SHALL BE ASSUMED BY THE PUTCHASER AND USER.

HELENA WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE PURPOSE REFERRED TO IN THE DIRECTIONS FOR USE SUBJECT TO THE INHERENT RISKS REFERRED TO ABOVE. HELENA MAKES NO OTHER WARRANTY OR REPRESENTATION OF ANY KIND, EXPRESS OR IMPLIED, CONCERNING THE PRODUCT, INCLUDING NO EMPLIED WARRANTY OF MERCHANTIBILITY OR FILMESS FOR ANY PARTICULAR PURPOSE, AND NO SUCH WARRANTLES SHALL BE IMPLIED BY LAW.

THE EXCLUSIVE REMEDY AGAINST HELENA FOR ANY CAUSE OF ACTION RELATING TO THE HANDLING OR USE OF THIS PRODUCT IS A CLAIM FOR DAMAGE AND IN NO EVENT SHALL DAMAGES OR ANY CHEER PRODUCT WITH AGAINST HELENA EXCEED THE PRICE OF THE PRODUCT WHICH CAUSES THE ALLEGED LOSS, DAMAGE, INJURY, OR OTHER CLAIM. HELENA SHALL NOT BE LIABLE AND ANY AND ALL CLAIMS AGAINST HELENEA ARE WAIVED, FOR SPECIAL, INDIRECT, INCIDENTAL, OR CONSEQUENTIAL DAMAGES OR EXPENSES, OF ANY NATURE, INCLUDING, EUT NO LIMITED TO, LOSS OF PROFITS OR INCOME, AND CROP OP PROPERTY LOSS OR DAMAGE, WHETHER OR NOT BASED ON HELENA'S NEGLIGENCE, BREACH OF WE CANTY, STRICT LIABILITY IN TORT OR ANY OTHER CAUSE OF ACTION.

HELENA AND THE SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPT IT, SUBJECT TO THE FOREGOING CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY, LIABILITY AND REMEDIES WHICH MAY ONLY BE VARIED BY AGREEMENT IN WRITING SIGNED BY A DULY AUTHORIZED REPRESENTATIVE OF HELENA.

PM12 60F6

.

ESTABLISHMENT NUMBER STATEMENT

ť

 \star First letters of product batch code indicate producing establishment.

5905-FL-1=TF · 5905-GA-1=CG · 5905-AR-1=WA 5905-OK-1=AO ·)905-TX-2=AT · 5905-CA-1=KC 5905-LA-3=DL