

CAUTION

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe dust. Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not apply when runoff is likely to occur.

Apply this product only as specified on this label. Dispose of wastes and containers by burying on non-crop lands away from water supplies.

ANTIDOTE

INTERNAL--Give sodium chloride 15 g. (one tablespoonful of table salt) in a glass of warm water and repeat until vomit is clear. Call a doctor immediately.

EXTERNAL—Wash with soap and water.

HELENA BRAND

15% GRANULAR INSECTICIDE

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED
FOR EUROPEAN CORN BORER CONTROL.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL**

ACTIVE INGREDIENT:	PERCENT BY WEIGHT
Toxaphene (technical chlorinated camphene containing 67-69%	15%
INERT INGREDIENTS	85%
TOTAL	100%

DIRECTIONS

EUROPEAN CORN BORER
13 pounds per acre
by airplane. A second application
7 to 10 days later.
Required for control of European Corn Borer.
State Extension Service for
application dates.

Do not ensilage material from
dairy animals or animals
intended for slaughter.

Do not allow dairy animals
to graze in treated areas
for slaughter to grade.

WARRANTY AND LIMITATIONS
AGES. Helena Chemical Company
this material conforms with the
on the label and is not warranted
referred to in the Directions
of Sale, subject to the terms
of the Helena Chemical Company
express or implied warranty
express or implied warranty
MERCHANTS AND DEALERS

HELENA

BRAND

15% GRANULAR INSECTICIDE

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED
FOR EUROPEAN CORN BORER CONTROL.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL**

ACTIVE INGREDIENT:	PERCENT BY WEIGHT
Toxaphene (technical chlorinated camphene containing 67-69%	15%
INERT INGREDIENTS	85%
TOTAL	100%

DIRECTIONS FOR USE

EUROPEAN CORN BORER CONTROL— Apply 13 pounds per acre by either ground equipment or by airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second-brood borers. Consult State Extension Service specialists for most effective application dates.

Do not ensilage made from treated corn plants to dairy animals or animals being finished for slaughter.

Do not allow dairy animals or animals being finished for slaughter to graze or feed on treated stover.

5-17-72
5905-186

WARRANTY AND LIMITATIONS OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of