

DANGER ← Red

DANGER: Poisonous by skin contact, inhalation, or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray. Wear clean heavy rubber boots, and clean water-proof protective clothing (coveralls, cap) and goggles when applying or handling. Destroy and replace gloves frequently. Wear effective mask or respirator of a type passed by the U. S. Bureau of Mines for EPN protection. Do not contaminate feed or foodstuffs. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water: for eyes, flush with water for 15 minutes and get medical attention.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

ANTIDOTES: Atropine is an antidote for EPN. Consult physician for emergency supply. Call a physician at once in all cases of suspected poisoning. Never give anything by mouth to an unconscious person. 2-PAM is also antidotal and may be administered in conjunction with atropine.

INTERNAL: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

EXTERNAL: Immediately remove all contaminated clothing and wash skin thoroughly with soap and warm water.

NOTE: Keep animals and unprotected persons away during treatment and while there is danger from drift or residues. Promptly destroy empty containers by burning or burying; stay out of smoke and vapors. Bury spillage; clean up area with strong lye solution.




6 TOX - 2 EPN

DANGER POISON KEEP OUT OF REACH OF CHILDREN SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

ACCEPTED

ACTIVE INGREDIENTS:	
*Toxaphene	53.45%
**O-Ethyl O-p-nitrophenyl phenyl- phosphonothioate	17.82%
Xylene	15.57%
INERT INGREDIENTS:	13.16%
TOTAL	100.00%

*Technical Chlorinated Camphene (67-69% Chlorine)
**Also known as EPN, U. S. Patent 2,503,390



DIRECTIONS FOR USE: Apply water to thoroughly cover plant surface or aircraft (1 to 3 gallons water by ground).

RECOMMENDATIONS FOR USE

BEANS (dry): Use 2-3 pints BEAN BEETLE, ARMYWORM (striped), THRIPS, SERPENTINE TED MITE, CLIMBING CUT (more than 21 days on beans. Do not feed animals being finished for slaughter.

CORN: Use 2 pints per acre to control TWO-SPOTTED MITES, MINER, ARMYWORMS, CLIMBING WORMS. Do not apply later than your State Agricultural Experiment Station treatments. Do not feed animals being finished for slaughter.

COTTON: Use 2-3 pints per acre to control APHIDS, THRIPS, BEANHOPPERS, TARNISHED PLANT BUGS, GARDEN WEBWORM. Do not feed animals being finished for slaughter in field; do not use within 3 days of harvest.


TOMATOES: Use 2-3 pints per acre to control TOMATO FRUITWORM, THRIPS, SERPENTINE LEAFMINER, TOMATO HORNWORM, and PINWORM. Do not apply later than 21 days before harvest.


Workers entering fields within 24 hours after application should wear protective clothing. Vacated areas should be treated with drifting insecticide and volatile residue.

WARRANTY AND LIMITATION: Helena Chemical Company warrants that the chemical described on the label conforms to the description and chemical description on the label for purposes referred to in the Directories, subject to the inherent risk of use. Helena Chemical Company makes no other warranty, including any other express or implied warranty of MERCHANTABILITY, and the Company is authorized to do so without reference to this warranty. Any other reference to this warranty shall be limited to include consequential commercial or value.



6 TOX - 2 EPN


DANGER POISON  **KEEP OUT OF REACH OF CHILDREN**
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL



ACTIVE INGREDIENTS:

*Toxaphene	53.45%
**O-Ethyl O-p-nitrophenyl phenyl- phosphonothioate	17.82%
Xylene	15.57%
INERT INGREDIENTS:	13.16%
TOTAL	100.00%

*Technical Chlorinated Camphene (67-69% Chlorine)
 **Also known as EPN, U. S. Patent 2,503,390



DIRECTIONS FOR USE: Apply specified rate per acre in sufficient water to thoroughly cover plant surfaces with ground driven equipment or aircraft (1 to 3 gallons water for aerial and 5 to 10 gallons water by ground).

RECOMMENDATIONS FOR USE

BEANS (dry): Use 2.2-3 pints per acre to control MEXICAN BEAN BEETLE, ARMYWORMS (beet, southern and yellow striped), THRIPS, SERPENTINE LEAF MINER, TWO-SPOTTED MITE, CLIMBING CUTWORMS. Do not apply later than 21 days on beans. Do not feed treated vines to dairy animals or animals being finished for slaughter.

CORN: Use 2 pints per acre to control EUROPEAN CORN BORER, TWO-SPOTTED MITES, THRIPS, SERPENTINE LEAF MINER, ARMYWORMS, CLIMBING AND SURFACE CUTWORMS. Do not apply later than 14 days before harvest. Consult your State Agricultural Experiment Station for proper timing of treatments. Do not feed treated forage to dairy animals or animals being finished for slaughter.

COTTON: Use 2.2-3 pints per acre to control BOLLWEEVIL, APHIDS, THRIPS, FLEAHOPPERS, RAPID PLANT BUGS, TARNISHED PLANT BUGS, COTTON LEAFWORM and GARDEN WEBWORM. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. Do not use within 3 days of harvest.

TOMATOES: Use 2.2-3 pints per acre to control ARMYWORMS, TOMATO FRUITWORM, FLEA BEETLE, APHIDS, THRIPS, SERPENTINE LEAFMINER, RUSSET MITE, TOMATO HORNWORM and PINEWORM. Do not apply later than 21 days before harvest.

Workers entering fields within 24 hours of treatment should wear protective clothing. Vacated areas should not be re entered until drifting insecticide and volatile residues have dissipated.

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

EPA REG. NO. 50075-101

DO NOT REUSE THE CONTAINER FOR ANY PURPOSE