WARNING

WARNING: Harmful if swallowed, inhaled, or absorbed through skin.

Keep out of reach of children, poultry, and domestic animals. Avoid breathing dust. Avoid contact with skin, wash thoroughly after using. Do not use inside homes or where milk is stored or processed. This product contains an organic phosphate which is a cholinesterase inhibitor and if poisoning should OCCUT CALL A PHYSICIAN IMMEDIATELY!

Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on label. Birds feeding on treated areas may be killed.


Do not contaminate feed, water or foodstuffs, milk or milking utensils. Not for use or storage in or around the home.

ATROPINE is the emergency antidote.

DIRECTIONS FOR USE

FLY BAIT may be used wet or dry. Kills flies fast in seconds. Easy to use. Simply walk and scatter lightly.

FLY BAIT is especially formulated to kill FLIES FAST and to provide longer residual fly control. Flies die within seconds after contacting or feeding on a particle of bait. The same bait may kill many flies before it loses its effectiveness. Treat only small areas where flies are a problem.



TO ATT WHEN CO A BINTS

(Contains Vapona® and Ronnel)

KILL FLIES IN SECONDS

KEEP OUT OF REACH OF CHILDREN. **WARNING:** See Other Precautions On Side Panel.

ACTIVE INGREDIENTS:

*2 2 dichlorovinyl dimethyl phosphate 0.093° *Related Compounds 0.007° Ronnel (O.O-dimethyl O-(2.4.5-trichlorophenyl)phosphorothicate 0.250°° **INERT INGREDIENTS** 99.650°-TOTAL 100.000°

*Vapona is a registered trademark of Shell Chemical Company

MANUFACTURED BY HELENA CHEMICAL COMPANY MEMPHIS. TENNESSEE

DIRECTIONS

PATIOS, DOG RUNS — Outisde use only. Scatter FLY BAIT lightly on surface areas where flies congregate. Use at the rate of 4 oz. (1/4 lb.) per 1500 square feet. Can be used on dry or slightly maistened surfaces. Repeat as necessary. Repeated use will help to keep the fly population below nuisance level.

GARBAGE CANS — Apply lightly around and inside garbage can. Repeat as

RESTAURANTS, DRIVE-INS, ICE CREAM STANDS, DOG KENNELS, OUT-DOOR MANTRINES, COMPOST AREAS, FOOD PROCESSING PLANTS, CANNERIES, EATING REAS -- Outdoor use. Sprinkle FLY BAIT lightly on floors, walks, allev-ways and ther horizontal surfaces where flies frequent. Repeat as necessary.

UNDER 1997 (N. I. A., AMAIN, TICIDE FUNGICION AND INTERIOR ECONOMIC FOREIGN REGISTER. ED UNDER NOSSESSION SUBJECT

DAIRY BARNS, STABLES, LIVESTOCK BARNS — Clean out manure and apply FLY BAIT on clean floor area or empty buriap sack in areas frequented by flies. Treat breas outside barns where flies cangregate. Use 4 oz. (% lb.) per 1000 square leet of area to be treated and repeat as necessary to reduce fly population.

CAGE LAYING OPERATIONS: KEEP OUT OF REACH OF FOULTRY — Scotter FLY BAIT on droppings under cages, on walkways and alley-ways. For reducing flies in and around Garbage Collection and Disposal Areas and Refuse Areas: Scatter lightly on ground, cement or other areas where flies congregate. "Iso at the rate of 2 oz. (1 8 lb.) per 1.000 square feet

LIQUID BAIT: Mix 1 pound of FLY BAIT in one gallon of water and sprinkle areas where flies congregate. Use at the rate of 1 pt. per 1,000 square feet.

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use and Conditions of Sale subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profit or value etc.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

NET WEIGHT

EPA Reg. No. 5905-221