

HELENA

BRAND

PAGE

IMPROVED GRANULES FOR SOIL INSECT CONTROL

ACTIVE INGREDIENTS:
 Technical Chlordane* 33.3%

INERT INGREDIENTS: 66.7%

Total 100.0%

*Equivalent to 19.98% Octachloro-4, 7-methanotetrahydroindane and 13.32% related compounds.

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE ANTIDOTE AND ADDITIONAL
PRECAUTIONARY STATEMENTS BELOW.

CAUTION: ← Red

Keep out of reach of children. Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing of dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE: ← Red

EXTERNAL - In case of spillage on skin, wash promptly with soap and water.

EXTERNAL - In case of spillage on skin, wash promptly with soap and water.

INTERNAL - Emetic of 1 tablespoonful of mustard in a tumbler of water. Call a physician.

Birds feeding on treated areas may be killed. This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other expressed or implied warranty, including any other expressed or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

CONTAINER DISPOSAL:
Do not reuse empty container.

Destroy container by burying with ~~soil~~ or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

Where row treatment is desirable, application should be made at planting time using suitable row equipment.

ACCEPTED

AUG 26 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5905-607

DIRECTIONS FOR USE
CROP OR USE

INSECT

PAGE
LBS ACRE
BROADCAST

LBS ACRE
TECHNICAL
CHLORDANE

PAGE
LBS ACRE
ROW

LBS ACRE
TECHNICAL
CHLORDANE

RECOMMENDATIONS AND LIMITATIONS

Keep out of reach of children. Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing of dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE: ← Red

EXTERNAL - In case of spillage on skin, wash promptly with soap and water.

EXTERNAL - In case of spillage on skin, wash promptly with soap and water.

INTERNAL - Emetic of 1 tablespoonful of mustard in a tumbler of water. Call a physician.

Birds feeding on treated areas may be killed. This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other expressed or implied warranty, including any other expressed or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

CONTAINER DISPOSAL:
Do not reuse empty container. ← Red

Destroy container by burying with roots or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

Where row treatment is desirable, application should be made at planting time using suitable row equipment.

ACCEPTED
AUG 26 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5905-227

DIRECTIONS FOR USE CROP (OR USE)	INSECT	PACE LBS./ACRE BROADCAST	LBS./ACRE TECHNICAL CHLORDANE	PACE LBS./ACRE ROW	LBS./ACRE TECHNICAL CHLORDANE	RECOMMENDATIONS AND LIMITATIONS
Vegetable Crops: Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cucumbers, Eggplants, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimentos, Potatoes, Radishes, Rutabagas, Squash, Sweet Potatoes, Tomatoes, Turnips, Watermelons.	White Grubs (Michigan only)	7½-12	2½-4	6	2	Do not make foliar applications to vegetable crops. On vegetable crops, make preplant broadcast or row application (where applicable) to soil after plowing and disk or incorporate into soil immediately. Do not graze treated areas to dairy animals or animals being finished for slaughter. Where a range of rates is given, use the higher rate of Helena Brand Pace on muck or highly organic soil.
	Whitefringed Beetle Larvae. (AL FL GA LA MS NC SC TN Only)	7½-15	2½-5			
Fruit & Berry Crops: Apples, Apricots, Cherries, Nectarines, Pears, Plums, Prunes, Quinces, Blackberries, Grapes, Huckleberries, Loganberries, Raspberries, Strawberries.	White Grubs (Michigan only)	7½-12	2½-4			On berry crops - Make application at base of plant and incorporate into soil. Do not apply when fruit is present. On tree fruits make soil application only when no fruit is present. Do not make foliar application to fruit and berry crops. Do not graze treated areas to dairy animals or animals being finished for slaughter. Where a range of rates is given, use the higher rate of Helena Brand Pace on muck or highly organic soil.
	Whitefringed Beetle Larvae. (AL FL GA LA MS NC SC TN Only)	7½-15	2½-5			
Strawberries. Pre-Plant treatment only.	Root Weevil Larvae	30	10			
	Strawberry Rootworm	15-30	5-10			
Non crop lands	Fire Ants	15-21	5-7			Do not use treated straw for feed or bedding. Do not graze treated areas or feed treated forage to dairy animals or animals being finished for slaughter. Where a range of rates is given, use the higher rate of Helena Brand Pace on muck or highly organic soil.

NET WEIGHT: 50 LBS.



EPA Reg. No. 5905-227