

**WARNING!**

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash after handling. Do not contaminate feed and food.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not use this product on feed or forage to be fed to Dairy animals or to livestock being finished for slaughter.

**DO NOT STORE NEAR HEAT OR OPEN FLAME.** Protect from temperatures below 0 degree F. Product crystallizes at lower temperatures. Warm or store at higher temperatures and mix to redissolve crystals and uniformly before use. Not for use or storage in or around the home.

Observe the following use recommendations. Read entire label before using. Use only for crops listed and only as specified on this label. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others. Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing, number of applications needed will vary with local conditions.

**DIRECTIONS FOR USE**

**DILUTE APPLICATION**  
Ground Application: Apply specified rate in 20 to 60 gal. of water per acre.  
Orchard Application: Apply specified rate in 100 to 800 gal. of water per acre.

**CONCENTRATE APPLICATION**  
Ground Application: Apply specified rate in not less than 5 gal. of water per acre.  
Orchard Application: Apply specified rate in 20 to 60 gal. of water per acre.

Special concentrate equipment is necessary for these uses.

Air Application: Apply specified rate in 5 to 20 gal. of water per acre. Orchard rates should not be applied in less than 10 gal. of water per acre.

**ALFALFA:** Alfalfa Caterpillar, Armyworms, Leafhopper, Alfalfa Looper, Clover Leaf Weevil, Corn Earworm, Climbing Cutworm, Grasshopper, Lygus Bug, Seed alfalfa Thrips, seed alfalfa Webworm. Apply 2 1/2 to 3 pts. per acre. Pea Aphid. Apply 3 pts. per acre. Apply in early spring or after cutting before new growth is 4 inches tall at the 2 1/2 pt. rate. Rates over 2 1/2 pts. or 3 pts. per acre do not feed treated forage to dairy animals being finished for slaughter.

**APPLES:** Red Banded Leafroller. Apply 6 to 16 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**BANANAS:** Leaf Eating Caterpillar, Cera M.D. Sees. Apply 1 pt. per acre. Do not apply within 1 day of harvest. Do not apply to treated areas to dairy animals or animals being finished for slaughter.

**BARLEY OATS RYE WHEAT:** Armyworm, Cutworm, and Sulfur Feeding Grasshoppers. Apply 2 1/2 to 3 pts. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use on the seed of these grains.

**BEANS:** Green Lima Snap. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray. Do not apply to treated areas to dairy animals or animals being finished for slaughter.

**BROCCOLI BRUSSEL SPROUTS CAULIFLOWER KOHLRABI:** Cabbage Looper, Cabbage Root Fly, Cabbage Root Weevil, Cabbage Worm, Cabbage Flea Beetle, Cabbage Aphid, Cabbage Thrips, Cabbage Flea Beetle, Cabbage Root Fly, Cabbage Root Weevil, Cabbage Worm, Cabbage Flea Beetle, Cabbage Aphid, Cabbage Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CABBAGE COLLARD KALE:** Cabbage Looper, Cabbage Root Fly, Cabbage Root Weevil, Cabbage Worm, Cabbage Flea Beetle, Cabbage Aphid, Cabbage Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CARROTS:** Cabbage Root Fly, Cabbage Root Weevil, Cabbage Worm, Cabbage Flea Beetle, Cabbage Aphid, Cabbage Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CELERY:** Celery Root Fly, Celery Root Weevil, Celery Worm, Celery Flea Beetle, Celery Aphid, Celery Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CITRUS:** Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines. Apply 6 to 16 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CLOVER:** seed crops only. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**CORN:** Corn Earworm, Corn Flea Beetle, Corn Rootworm, Corn Aphid, Corn Thrips, Corn Flea Beetle, Corn Rootworm, Corn Aphid, Corn Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**COTTON:** Armyworm, Bollworm, Boll Weevil, Cotton Flea Beetle, Cotton Rootworm, Cotton Aphid, Cotton Thrips, Cotton Flea Beetle, Cotton Rootworm, Cotton Aphid, Cotton Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**EGGPLANTS PEPPERS:** Eggplant Flea Beetle, Eggplant Rootworm, Eggplant Aphid, Eggplant Thrips, Eggplant Flea Beetle, Eggplant Rootworm, Eggplant Aphid, Eggplant Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**HICKORY NUTS:** Hickory Nut Weevil, Hickory Nut Flea Beetle, Hickory Nut Rootworm, Hickory Nut Aphid, Hickory Nut Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**HORSERADISH:** Horseradish Flea Beetle, Horseradish Rootworm, Horseradish Aphid, Horseradish Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**LETTUCE:** Lettuce Rootworm, Lettuce Flea Beetle, Lettuce Aphid, Lettuce Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**OKRA:** Okra Flea Beetle, Okra Rootworm, Okra Aphid, Okra Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

**ONIONS:** Onion Flea Beetle, Onion Rootworm, Onion Aphid, Onion Thrips. Apply 2 1/2 to 3 pts. per acre. Do not apply within 40 days of harvest or after seed and tree spray.

# HELENA

## BRAND

### TOXAPHENE

#### SPRAY 8-E

This product contains 8 pounds of Toxaphene per gallon.

**WARNING** KEEP OUT OF REACH OF CHILDREN  
SEE OTHER WARNINGS ON SIDE PANEL

ACTIVE INGREDIENTS	8.00
INERT INGREDIENTS	12.00
TOTAL	20.00

BEST AVAILABLE

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE

MANUFACTURED BY  
**HELENA CHEMICAL COMPANY**  
MEMPHIS TENNESSEE

PARSNIPS  
PEACHES  
PEAS  
PIMENTOS  
PINEAPPLE  
QUINCE  
RICE  
RUTABAC  
SORGHUM  
SOYBEAN  
SPINACH  
STRAWBERRY  
TOMATO  
TURF SEED  
ORNAM