

CAUTION!

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF INHALED OR SWALLOWED
AVOID BREATHING OF SPRAY
DO NOT TAKE INTERNALLY
AVOID CONTACT WITH SKIN AND EYES
WEAR REGULAR LONG-SLEEVED WORK CLOTHING
CHANGE TO CLEAN CLOTHES DAILY
WASH HANDS AND FACE BEFORE EATING
WASH THOROUGHLY AFTER HANDLING

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do Not Use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS During Application Or When Cleaning Equipment

NOT TO BE STORED IN AREAS WHERE TEMPERATURES FREQUENTLY EXCEED 100°F.

GENERAL CAUTIONS

- 1) Physical compatibility of SEVIMOL with other pesticides is not fully known. Incompatible mixtures curdle or precipitate and may injure crops or reduce control. Compatibility can be affected by water in your locality, various surfactants and other formulation ingredients. Do not apply if physically incompatible.
- 2) Do not mix SEVIMOL with nutrients or adjuvants or other pesticides unless your experience indicates that the mixture is safe and effective. If uncertain, prepare a small amount of the mixture and test on a representative portion of the crop.
- 3) Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime-sulfur and casein-lime spreaders.
- 4) To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

BEE CAUTION: May kill honeybees in substantial numbers.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use when value of bees as pollinators is more important than insect control. Before applying warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

SEVIMOL® is a suspension of SEVIN® carbaryl insecticide in molasses for dispersion in water and low volume application in hydraulic sprayers, mist blowers, low gallonage ground equipment, and aircraft. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when SEVIMOL is used in accordance with label directions. If SEVIMOL is used in accordance with label directions, forage or feed crops may be used or harvested for use as feed for dairy or meat animals without resulting residues in meat or milk. **READ THIS LABEL. USE IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS.**

DIRECTIONS FOR EQUIPMENT CARE, SPRAY PREPARATION AND CONTAINER DISPOSAL

Equipment Care:

- 1) Flush entire spray system with clear water with clear water numerous times before using SEVIMOL. **DO NOT FLUSH WITH KEROSENE OR DIESEL FUEL.**
- 2) Use 50 mesh or coarser screens in entire system.
- 3) Nozzles should be fan type, number 3 or larger. Cone nozzles may be used if applying at least 3 gallons of finished spray per acre.
- 4) After spraying, flush tank and lines with clear water.

Spray Preparation:

- 1) SEVIMOL may be applied undiluted on cotton. For other crops, see directions 2 through 5.
- 2) Add required amount of water to spray or mix tank and provide moderate agitation while adding SEVIMOL. Refer to specific crop headings for directions on amount of water to use.

3) For combination sprays SEVIMOL must be added to at least an equal volume of water and mixed thoroughly before another pesticide is added. (See **General Cautions**).

4) Recirculate spray mixtures at frequent intervals after mixing to maintain a uniform suspension.

5) Mix only as much spray as can be applied on day of mixing.

Storage of Unused Material and Container Disposal:

- 1) Store unused SEVIMOL in original container only, in cool, dry area out of reach of children and animals.
- 2) Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

IMPORTANT: The directions on this label are based on tests and field experience relating to (a) effectiveness; and (b) possible injury to plants and animals; and (c) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

COTTON INSECT CONTROL

Recommended dosages refer to quarts of SEVIMOL per acre. Molasses in formulation intended for improved bollworm control.

Use ½ to 1 quart for early season Thrips, Flea Beetles, Cotton Fleahopper, Striped Blister Beetle and Cotton Leafworm.

Use 1 to 2 quarts, depending on the size of the cotton and insect population level, for mid- and late-season Boll Weevil, Bollworms, Fall Armyworms, Cotton Leaf Perforator, Leaf Rollers, Leafhoppers, and Tarnished Plant Bug. Also for light-to-moderate infestations of Lygus Bugs in Western irrigated cotton. Applications should be made every 5 to 7 days as long as control is necessary.

Use 1½ to 2½ quarts for Pink Bollworm on a 5 to 7 day schedule depending on the size of the cotton and degree of infestation.

Use 2 quarts for Stink Bug and Salt-Marsh Caterpillar.

Use ½ to 1½ quarts for Grasshoppers. The highest dosage should be used when Grasshoppers are mature or material is applied on partly defoliated plants.

Aphid populations will be suppressed by repeated application of this insecticide. SEVIMOL does not control Spider Mites. Where infestations are encountered, use a recommended miticide.

To avoid staining lint, do not use SEVIMOL after bolls open.

FORAGE AND FIELD CROP INSECT CONTROL

Recommended dosages refer to quarts of SEVIMOL per acre. Apply when insects or their damage appear and repeat 7 to 14 days later if necessary. Apply sufficient gallonage to obtain full coverage but not less than one gallon of finished spray per acre. See specific directions below for Grasshopper control.

ALFALFA AND SOYBEANS: Recommended dosages refer to quarts of SEVIMOL per acre.

Use ½ to 1 quart for Blue Beetle and Mexican Bean Beetle.

Use 1 quart for Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Leafhoppers, Three-Cornered Alfalfa Hopper, Thrips, Velvet Bean Caterpillar, Green Cloverworm and Japanese Beetle.

Use 1 to 1½ quarts for Armyworms, Corn Earworm (Bollworm), Stink Bugs and Webworms.

Use 1½ quarts for control of Alfalfa Weevil larvae in the spring. If pretreatment damage is extensive, cut the alfalfa and apply to stubble.

Use 1½ quarts for Cutworm control on alfalfa.

To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

WARNING! Do not apply in combination of SEVIMOL and 2,4-DB herbicide (such as "Butyrac" 75 or "Butoxone" 5B) to soybeans.

SORGHUMS (Including MILO and GRAIN SORGHUM): Use 1 to 2 quarts for Armyworms, Corn Earworm, Stink Bugs and Webworms. Use 1½ quarts for Sorghum Midge and Southwestern Corn Borers.

(Directions for Use continued on right panel)

STOP — READ LABEL BEFORE USING!



SEVIMOL 4

Carbaryl Insecticide

This Product Contains 4 Pounds Carbaryl Per Gallon

For Control of Insect Pests
Especially for Improved Bollworm Control

CAUTION Keep Out of Reach
of Children. See
Other Caution Statements on Side Panel

SEVIMOL® is the TM of Union Carbide Corporation for an insecticide containing Sevin® carbaryl and molasses.
U.S. Pat. Nos. 2,903,478 and 3,009,855

Active Ingredient:	By Wt.
Carbaryl (1-naphthyl methylcarbamate).....	40.38 %
Inert Ingredients:	59.62 %
Total	100.00 %

**DO NOT RE-USE THIS CONTAINER
FOR ANY PURPOSE**

MANUFACTURED BY
HELENA CHEMICAL COMPANY
MEMPHIS, TENNESSEE

5905-251

1-5-1977

2/1

SORGHUMS (Continued):

Use 2 quarts for Cutworm control. For optimum insect control on grain sorghum, direct spray into the forming heads. Treat for Sorghum Midge 3 to 4 days after heads have emerged from boot.

Do not apply within 21 days of harvest of grain. No time limitation on sorghums used for forage.

SUGAR BEETS: Use 1 to 2 quarts for Armyworms, Leafhoppers, and Webworms.

Do not apply within 14 days of harvest.

GRASSHOPPERS: Apply 1/2 to 1 1/2 quarts as often as necessary to control Grasshoppers on the above forage crops. Follow pre-harvest and grazing use limitations listed for each of the above crops. A 1/2 to 1 quart rate is suggested for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditchbanks, and borders.

A 1 to 1 1/2 quart dosage should be used when Grasshoppers are mature or material is applied to crops requiring greater coverage.

TOBACCO INSECT CONTROL

Recommended dosages refer to quarts of **SEVIMOL 4** per acre.

Special Instructions: When using conventional ground equipment, dilute with water and use 25-40 gallons of finished spray per acre. Apply for full coverage.

Use 1 to 2 quarts for Budworm, Flea Beetle, Japanese Beetle, June Beetle and Hornworm. Use lower rate on young plants up to knee high; higher rate for mature plants. Begin treatment when worms are small. Avoid excessive application in plant bud as injury may result.

Treated fields may be entered immediately after foliage has dried.

If late applications are necessary, allow 3 days before painting or cutting.

SMALL FRUIT INSECT CONTROL

Recommended dosages refer to quarts of **SEVIMOL** per acre.

Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7- to 10-day intervals or as necessary.

CRANBERRIES: Use 2 quarts for Cutworms, Fireworms, Fruitworms, Japanese Beetle and Leafhopper. Apply in late bloom and as needed at 7- to 10-day intervals. A dilute spray of 300 gallons per acre is suggested.

Do not apply within 1 day of harvest.

STRAWBERRIES: Use 1 to 2 quarts for Strawberry Leaf Roller. A dilute spray of 100 to 200 gallons per acre is suggested.

Do not apply within 1 day of harvest.

VEGETABLE INSECT CONTROL

Suggested dosages refer to quarts of **SEVIMOL** per acre. Where a dosage range is indicated, use lower rate on young plants; higher on mature plants.

Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7- to 10-day intervals or as necessary unless shorter interval is specified below. Apply sufficient gallonage to obtain full coverage but not less than one gallon of finished spray per acre. See specific directions below for Grasshopper control.

CORN: Use 1 1/2 quarts for Corn Earworm, Corn Rootworm adults, Grasshoppers, European Corn Borer, Southwestern Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetle and Leafhopper. Use 2 quarts for Western Bean Cutworm. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at least every 2 or 3 days starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation; timing and good coverage are absolutely essential for effective control.

Application of **SEVIMOL** to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Use 2 quarts in at least 15 gallons of water for Cutworm control. Spray in 17 inch band over corn row.

BEANS: (Green Beans, Lima Beans, Snap Beans, Cowpeas, and Black-Eyed Peas).

Use 1/2 quart for Mexican Bean Beetle.

Use 1 quart for Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvet Bean Caterpillar and Western Bean Cutworm.

Use 1 to 1 1/2 quarts for Armyworms, Cutworms, Corn Earworm, Stink Bugs, and Tarnished Plant Bug.

In California, use 2 quarts for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bugs.

POTATOES, TOMATOES, AND PEPPERS: Use 1/2 to 1 quart for Colorado Potato Beetle, Flea Beetles, and Leafhoppers. Use 1 to 2 quarts for European Corn Borer, Fall Armyworm, Flea Bugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug and Stink Bugs. Use 2 quarts for Cutworm control.

GRASSHOPPERS: Apply 1/2 to 1 1/2 quarts as often as necessary to control grasshoppers on the above vegetable crops.

SHADE TREE AND ORNAMENTAL INSECT CONTROL

Use 1 quart per 100 gallons of water (1 tablespoon per gallon) when insects or their damage appears. Repeat weekly or as needed.

Use 1 to 2 quarts per 100 gallons for control Eriophyid Mites. In dilute applications the addition of an emulsifiable horticultural oil may enhance the control of Eriophyid Mites. Carbaryl injures Boston Ivy and Virginia Creeper.

Use on herbaceous and annual plants (Carnation, Chrysanthemum, Gladiolus, Iris, Peony, Zinnia, etc.) to control:

Blister Beetle	Leafroller
Boxelder Bug	Mealy Bug
Flea Beetle	Plant Bug
Japanese Beetle	Psyllid
June Beetle	Rose Aphid
Lace Bug	Thrips (exposed)
Leafhopper	

Use on shrubs, trees and woody plants (Ash, Arborvitae, Azalea, Bayberry, Beech, Birch, Boxwood, Catalpa, Cedar, Cypress, Dogwood, Elm, Euonymus, Ginkgo, Hackberry, Hawthorn, Holly, Honey-suckle, Hydrangea, Juniper, Lilac, Magnolia, Maple, Oak, Pine, Redbud, Rose, Tulip-tree, etc.) to control:

Apple Aphid	Mealy Bug
Bugworm	Oak Leaf Miner
Black Leaf Miner	Orange-striped Oakworm
Boxelder Bug	Orange Tortix
Boxwood Leaf Miner	Periodical Cicada
Cankerworms	Pine Sawfly
Catalpa Sphinx*	Puss Caterpillar
Cooley Spruce Gull Aphid	Plant Bug
Eastern Spruce Gull Aphid*	Rose Aphid
Elm Leaf Aphid	Rose Slug
Elm Leaf Beetle	Scale Insects
Elm Spanworm*	Spruce Needle Miner*
Eriophyid Mite*	Tent Caterpillar
Gypsy Moth	Thorn Bug
Japanese Beetle	Thrips (exposed)
June Beetle	Webworms
Lace Bugs	Willow Leaf Beetle
Leafhopper	Yellow Poplar Weevil*
Leafroller	

IMPORTANT: Use clean equipment. Flush tank and lines with clean water before and after spraying.

Sprays should be 50 mesh or coarser.

Provide moderate agitation while adding **SEVIMOL** to water in spray or mix tank. Recalculate at intervals to maintain a uniform suspension.

WARRANTY AND LIMITATIONS OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profit or value, etc.

EPA Reg. No. 5905-25

ACCEPTED

JAN 5 1977

NET CONTENTS:

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

2/c