

卷之三

**KILL OUT OR EACH OF CHILDREN
HAPPYLY TO WAITED.**

HELENA

Avoid contact with sites, skin and Do not breathe dust.

Each thoroughfare after visiting.

AMONG THE TOWNSHIP MERCHANTS
AND COTTON DEALERS.

20% TOLUAPHEN -
40% SULPHUR JUST

An Agricultural Insecticide List

ACCEPTED

2 AUG 1974

CAUTION TO KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTION STATEMENTS ON
THIS PAGE.

בבב ביבריאן טראניון סאלומון סאלומון זייד פאבר

卷之三

卷之三

卷之三

卷之三

卷之三

卷之三

卷之三

卷之三

COTTON: Cotton is less open to wintering than to summering. The cotton is wintered in bolls which are present. The cotton is wintered in bolls which are present. The cotton is wintered in bolls which are present. The cotton is wintered in bolls which are present.

Leafworms, Fall Armyworms, Flea Beetles, Grasshoppers, Ticks and other Mirids - Use 10 to 15 lb. per acre. Apply as needed at 7 to 10 day intervals. Roll vegetables, earthworms, cutworms, Garden Webworms - use 15 to 20 lb. per acre. Apply at 4 to 5 day intervals until control is obtained. Use 20 lb. per acre.