



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

259579

Application for Pesticide - Section I

1. Company/Product Number 5905-302	2. EPA Product Manager A. Abrahamovich	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Sol Oil Plus	PM#	
5. Name and Address of Applicant (Include ZIP Code) Helena Chemical Company 6075 Poplar Avenue, Suite 500 Memphis, Tennessee 38119 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input checked="" type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

AUG 20 2001

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Amendment by Notification to Correct Error in First Aid Statement (previously amended per PR Notice 98-10) typographical errors exist and to amend warranty statement. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Emily Smathers Ratliff	Title Registration Supervisor	Telephone No. (Include Area Code) 901/752-4421
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Supervisor	
4. Typed Name Emily Smathers Ratliff	5. Date 08/02/01	

2/7

SOL OIL PLUS

INSECTICIDE-MITICIDE/SPRAY

ACTIVE INGREDIENT:

Petroleum oil 90.00%

INERT INGREDIENTS: 10.00%

TOTAL 100.00%

Petroleum Oil Min. Unsulfonated Residue - 96-98%

Protected by U.S. Patent # 5,462,912

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Other Cautions on Inside Panel

EPA REG. NO. 5905-302

EPA EST NO.

NET CONTENTS:

NOTIFICATION

AUG 20 2001

MANUFACTURED BY
HELENA CHEMICAL COMPANY
MEMPHIS, TN. 38119

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID	
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
If on Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on Skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>NOTE TO PHYSICIAN: This product contains petroleum distillate - vomiting may cause aspiration pneumonia.</p>	

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber or viton.
- Shoes plus socks
- Chemical -resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Wash hands before eating drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour spill or store near heat or open flame.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber or viton.
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses. Do not enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Keep container well sealed. Protect from excessive heat.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, by other procedures approved by state and local authorities.

SOL OIL PLUS is a special blend of refined phyto bland oils that function as a contact insecticide and miticide. In addition, the surfactant/emulsifier system utilized in SOL OIL PLUS improves spray application by modifying the wetting and deposition characteristics of the spray droplets.

For foliage application, the spray tank must be free of any sulfur residue before using SOL OIL PLUS. Do not combine sulfur sprays with SOL OIL PLUS on foliage and do not follow a previous application of sulfur or apply sulfur following SOL OIL PLUS until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulfur, using 1 quart to 25 gallons water. If sufficient time has not elapsed, burn will shown in several days. When temperatures drop below 32o F (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35o F and are free from moisture due to melting frost or heavy dews. SOL OIL PLUS can be used with recommended rates of other pesticides. Application to trees weakened by disease, drought, driving winds or high nitrogen applications may result in oil injury.

MIXING INSTRUCTIONS

- 1) Add enough water to the mixing tank for proper agitation.
- 2) Add nutrients and other pesticides. If dry pesticide products are used, mix the materials with water thoroughly so that they are completely dispersed in the mixing tank. DO NOT add SOL OIL PLUS before a uniform mixture is obtained.
- 3) Maintain agitation of the mix until the spray is applied. Flush any spray mix remaining in hoses back into the tank if the solution stands for more than 20 minutes.
- 4) Ensure that spray equipment is thoroughly clean so that residues from other applications do not remain.

For all applications to be effective, complete coverage is essential, necessitating a thorough inside and outside tree application.

ALFALFA (Seed Crop Only): Use 1 quart per acre in a minimum of 20 to 30 gallons of water per acre by ground application or a minimum of 10 to 15 gallons by air when used in combination and at the recommended label rates with insecticides or miticides that recommend the use of oil. To increase the effectiveness of desiccants, apply 2 gallons per

acre in a minimum of 10 gallons of spray by air. Use is limited to those herbicides/desiccants labeled for Alfalfa Seed desiccation. Spider Mites: Use 1 quart per acre at the dilutions noted above.

ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES - Dormant or Delayed Dormant: Dilute spray: Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale (Parlatoria Scale), Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Parlatoria, Soft Brown Scale, other Scale Insects, Eggs of Aphids, Red Spider Mites, Brown Mites - 1½ to 2 gallons in 300 to 600 gallons per acre. Concentrate spray: 4 to 6 gallons per acre in a minimum of 20 gallons of spray. Almond and Apricots: for the European Red Mite, Brown Mite, San Jose Scale - 4 to 8 gallons per acre in a minimum of 20 gallons of spray. Spring or Summer Dilute. Spray: Twospotted, Pacific, European Red and Brown Mites: 1 to 2 gallons in 300 to 600 gallons of water per acre. Spring or Summer Concentrate: 4 to 6 gallons per acre in a minimum of 20 gallons of spray. Application by air: Use no more than 4-6 gallons of product in a minimum of 20 gallons of spray. Plums and Prunes: for Peach Twig Borer - 1½ - 2 gallons per 100 gallons of water. Concentrate spray: 6 - 8 gallons per 20 - 125 gallons of water. Application by air: 6 - 8 gallons in a minimum of 20 gallons of spray.

APPLES: Rust Mite: Delayed Dormant - 1½ to 2 gallons. Summer use - 1 to 1½ gallons.

APPLES, PEARS - Dormant or Delayed Dormant - Dilute Spray: San Jose Scale, Pear Scale, Eggs of Aphids, Red Spider Mites (Rust Mites, Apples only - delayed dormant) - 1½ to 2 gallons. Pears only: Delayed dormant. Pear Psylla - 2 gallons. Concentrate spray - use 4 to 6 gallons in a minimum of 20 gallons of spray per acre. Summer dilute spray: Twospotted Spider, Pacific and European Red Mites - 1 to 1½ gallons dilute in 400 to 800 gallons of water. Concentrate spray: use no more than 4 gallons per acre in a minimum of 20 gallons of spray. Application by air: use no more than 4 - 6 gallons of product in a minimum of 20 gallons of spray per acre. (Concentrated Sprayers - Pears (California Only) - 4 to 6 gallons in 90 to 120 gallons of water per acre).

Dormant: (Washington and Oregon only): Apples - Dilute spray for mites and San Jose scale - use 1 to 1½ gallons of oil plus sulfur or lime sulfur at the label recommended rate. Concentrate spray - use 4 to 6 gallons of oil per acre plus the label recommended rate of sulfur or lime sulfur in 20 to 100 gallons of spray per acre. Pears - Dilute Spray - Pear Psylla - use 1 to 1½ of oil plus sulfur or lime sulfur at the label recommended rate. Concentrate spray - use no more than five gallons of oil plus the label recommended rate of sulfur or lime sulfur in 20 to 100 gallons of spray per acre.

NOTE: Oil plus sulfur or lime sulfur can cause severe foliage and fruit bud injury if applied after the dormant period.

ASPARAGUS, BEANS, BEETS, CORN, CUCURBITS, PEPPERS, SQUASH, & TOMATOES: Aphids, Mites, Beetle Larvae, Leafminers, and Certain Caterpillars - Use 1 - 2 gallons/100 gallons of water.

AVOCADOS (HAAS VARIETY ONLY): Avocado Brown Mite - 1 - 1 1/2 gallons per 100 gallons with thorough coverage through late fall.

CITRUS (Citrus spp., Fortunella spp.) Florida Red Scale, Purple Scale Soft Brown Scale, Black Scale, White Flies, Red Spider Mites, Rust Mites - Concentrate spray: 2 gallons per acre if total spray volume is less than or equal to 125 gallons per acre. If total spray volume is greater than 125 gallons per acre, use a 2% volume for volume. Control of severe insect infestation may require a higher rate not to exceed 3 gallons per acre.

NOTE: Do not exceed 500 gallons per acre total spray volume. Do not use within 8 weeks of harvest on Lemons. Do not use on Navel Oranges after September 1.

ALL CITRUS VARIETIES: Do not exceed 0.4% v/v concentration. Do not use oil when freezing temperatures are expected within 24 hours of the application. Care should be taken in combining oil sprays with certain miticides and insecticides, due to possible increased phytotoxicity when applied over foliage. See an Agriculture Extension Service agent for recommendations on special combinations.

FIGS: Fig Scale - Dormant or Delayed Dormant: Dilute Spray - use 2 to 3 gallons per 100 gallons in 300 to 600 gallons per acre water. Use 3 gallons for severe infestations. Concentrate spray - use 4 to 6 gallons in a minimum of 50 gallons of spray.

GRAPES: Dormant : Dilute Spray: Mealybug - Use 1 to 2 gallons plus the appropriate labeled rate of an insecticide registered for use on grapes for mealybug control in 200 to 300 gallons water per acre. Concentrate spray - use 1 to 2 gallons plus the appropriate labeled rate of an insecticide registered for use on grapes for mealybug control per acre in enough water for thorough coverage. Suppression of Grape and Variegated leafhopper nymphs suppression - 1 to 1½ gallons or 1 - 3 gallons per acre concentrate. Mite suppression - 1 - 2 gallons per acre concentrate. Narrow range oils such as SOL OIL PLUS can temporarily affect surface bloom appearance but will not cause subsurface injury. When

applied for leafhopper or mite suppression near harvest (within 2 days) SOL OIL PLUS should not be applied to grapes for fresh market.

OLIVES: Dilute spray: Parlatoria, Oleander, Olive and Black Scales, Spider Mites - use 1½ in 400 to 500 gallons per acre water. May be used post bloom. (Late May through June when crawlers are present) Summer (July 15 through August) and Post-Harvest (October through November).

PEARS: Pear Psylla - Delayed Dormant - 2 gallons per acre. Summer Use - 1 to 1-1/2 gallons. Blister Mite - Delayed Dormant and Fall Spray - 2 gallons per acre.

PECANS: Obscure Scale - Dormant: Dilute Spray - use 3 gallons in 100 gallons water per acre. Concentrate spray - use 6 to 8 gallons per 20 - 125 gallons water per acre.

STRAWBERRIES: Twospotted Spider Mites - Use 1.0% to 2.0% volume to volume in 60 gallons of water/acre with air assisted low volume ground applicators or 300 gallons of water/acre with standard ground spray equipment. Application may be made during the winter months when plants are semi-dormant. Do not use oil from peak bloom through peak fruiting period. Application should be made when mites are at no more than a moderate infestation level. Higher levels of mite infestations will require treatment with more effective miticides. Make applications at no less than 14 days between applications. Consult your local Extension Office for more specific local mite threshold density recommendations for this type of treatment. When using refined petroleum oils on strawberries there will be a danger of phytotoxicity. To minimize the problem, do not apply this product when warm temperatures, 75 degrees F and higher, are expected for several days after application. In California, do not apply this product from:

January 16 to May 30 in Orange and San Diego counties and the Oxnard Plains
February 1 to June 15 in the Santa Maria Valley
March 1 to June 30 in Monterey and Santa Cruz counties.

SHADE TREES, SHRUBS, ORNAMENTALS, FLOWER AND FOLIAGE PLANTS, CHRISTMAS TREES

Shade Trees* and Shrubs including Conifers (See Note), Deciduous Broadleaf Evergreens & Woody Ornamentals: Aphids, Adelgids, Bugs (immature), Certain Caterpillar, Eriophyid Mites, Gall Mites, Lace Bug, Leaf Beetle, Larvae, Leafminer, Mealybug, Psyllids (immature), Sawfly (larvae), Scales (immature), Spider Mites, Whiteflies (immature): Winter Dormant Period as needed - 2 to 4 gallons oil per 100 gallons water. Summer (Foliar or Cover) as needed - 1 to 3 gallons oil per 100 gallons water. Note: May cause discoloration of Blue Spruce. Do not use on Conifers or Ferns not tolerant to oil sprays.

Ornamental Trees*, Shrubs along city streets, other rights-of-way including Conifers, Deciduous and Broadleaf Evergreens: Aphids, Adelgids, Bugs (immature), Certain Caterpillar, Eriophyid Mites, Gall Mites, Lace Bug, Leaf Beetle Larvae, Leafminer, Mealybug, Psyllids (immature), Sawfly (larvae), Scales (immature), Spider Mites, Whiteflies (immature): Winter Dormant Period as needed - 2 to 4 gallons oil per 100 gallons water. Summer (Foliar or Cover) as needed - 1 to 3 gallons oil per 100 gallons water.

Flowering and Foliage plants including Roses and other Flowering Shrubs: Aphids, Adelgids, Bugs (immature), Certain Caterpillar, Eriophyid Mites, Gall Mites, Lace Bug, Leaf Beetle Larvae, Leafminer, Mealybug, Psyllids (immature), Sawfly (larvae), Scales (immature), Spider Mites, Whiteflies (immature): Winter Dormant Period as needed - 2 to 3 gallons oil per 100 gallons water. Summer (Foliar or Cover) as needed - 1 to 2 gallons oil per 100 gallons water.
Foliage Ornamentals & Bedding Plants: Aphids, Adelgids, Bugs (immature), Certain Caterpillar, Eriophyid Mites, Gall Mites, Lace Bug, Leaf Beetle Larvae, Leafminer, Mealybug, Psyllids (immature), Sawfly (larvae), Scales (immature), Spider Mites, Whiteflies (immature), Winter Dormant Period as needed - 2 to 3 gallons oil per 100 gallons water. Summer (Foliar or Cover) as needed - 1 to 2 gallons oil per 100 gallons water.

Christmas Trees: Aphids, Adelgids, Scale (soft & hard) (immature), Spider Mites: 2 to 4 gallons oil per 100 gallons water. Summer 1 to 3 gallons oil per 100 gallons water.

*Oil removes the glaucous (blue) bloom from such evergreens as Colorado Blue Spruce and Koster Spruce. Always use lower dosage or test spray oil sensitive plants such as Cryptomeria, Smoke Tree, Chamaecypris, Juniper, Japanese Holly and Spruce. Tendency toward sensitivity: Red Cedar and Douglas Fir. Do not spray Walnut foliage. Do not use on Conifers or Firs not tolerant to oil sprays.

Note: Spray no more than 4 times during the growing season; use a 2 week minimum application interval. Do not spray when buds have fully opened and shoot elongation is occurring. Do not spray when there is obvious moisture deficit in leaves, or the plant is under stress. For fall dormant applications, reduce rate to 2 gallons oil per 100 gallons water and

limit use only to: American Red Oak, Japanese Black Pine, Dogwood, Weeping Cherry, Cornelian Cherry, Crabapple, Norway Maple, Purple Leaf Plum. Keep away from open blooms - bleaching and spotting has been observed with the open blooms of certain plants.

WALNUTS - Late Spring to Mid-Summer Only: Summer Stages of Frosted Scale, European Fruit Scale, San Jose and other Soft-Shell Scale, Red Spider Mites - 2 to 4 quarts. The addition of one of the following phosphate insecticides aids in scale control, Trithion or Ethion. See label for recommended dosages. Delayed dormant: Black, San Jose, Olive, Frosted, Calico, European Fruit Leucanium Scales, Red Spider Mite Eggs and Walnut Aphid Eggs - Dilute Spray - use 3 gallons in 300 to 500 gallons. Foliar (Fully expanded leaves): Spider Mites - use 4 gallons per acre in 200 to 400 gallons of water. Concentrate Spray: for delayed dormant and foliar sprays, apply in a minimum of 40 gallons of water per acre. Use oil only on trees that have not suffered from lack of moisture or other stressing factors during the growing season. Do not apply when soil moisture is low. Do not apply after husk split. As an adjuvant for neutral copper compounds for control of Walnut Blight in pre and post-bloom sprays - use 1 quart per 100 gallons of water as a dilute spray. Follow dosages and timings on the copper product label.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Chemical Company, 2001.
**Keltthane® is a Reg. Trademark of Rhom and Haas Company
SOL OIL PLUS is a trademark of Helena Chemical Company.

