

CAUTION
Keep Out of Reach of Children
HARMFUL IF INHALED OR SWALLOWED.

Do not breathe dust. Do not take internally.

Avoid prolonged or repeated breathing of dust. Skin contact may be harmful. Avoid contact.

Wash hands thoroughly after handling and before eating or smoking. Take shower or bath after work. Wear long sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7 to 10-day intervals when needed to maintain control. Observe use limits given for specified crops.

OBSERVE THESE PRECAUTIONS

Sevin Insecticide will not control insects if used with lime or other highly alkaline materials.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Do not contaminate feed or foodstuffs.

Avoid drift to food and freezer crops.

HELENA
BRAND
SEVIN* 5
DUST

Trade Mark of Union Carbide Corp. for the active ingredient carbaryl, 1-naphthyl N-methylcarbamate

U. S. Patent No. 2,903,478

CAUTION KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl, 1-naphthyl N-methylcarbamate	5%
INERT INGREDIENTS	95%
TOTAL	100%

Notice: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions and under normal use, but neither this warranty, nor any other warranty of merchantability or fitness for particular purpose, if expressed or implied, extends to the use of this product contrary to the label directions, or to the use of this product under conditions not expressly foreseeable to seller, and the user assumes the risk of any such use.

COMPLETELY EMPTY CONTAINER. BURY UNUSED CHEMICAL AT LEAST 18 INCHES DEEP IN AN ISOLATED LOCATION, WELL AWAY FROM WATER SUPPLIES. BURN EMPTY CONTAINER COMPLETELY. EXERCISE EXTREME CAUTION AND STAY WELL AWAY FROM SMOKE. BURY ASHES AND RESIDUE.

RECOMMENDATIONS

POULTRY (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite Lice and as a supplement to premise treatments for Chicken Mites, Fleas — 1 lb. per 100 birds. Apply thoroughly by means of a shaker or hand duster. Repeat in 4 weeks if necessary. Do not use on poultry or game birds within 7 days of slaughter.

IN OR AROUND POULTRY HOUSES: Chicken Mites, Fleas, Bed Bugs — 1 lb. per 40 sq. ft. of litter or floor space and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatments within 7 days of slaughter. Avoid contamination of feed, feeding troughs and watering receptacles.

CORN: Corn Earworms, Fall Armyworms, European Corn Borers — Use 30 lb. per acre starting when the silks first appear. Repeat at 2 to 3 day intervals throughout the silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. May be applied up to harvest on sweet corn ears for human consumption. To avoid excessive residues on fodder and forage, do not apply within 7 days of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

PEANUTS: Corn Earworms, Fall Armyworms, Potato Leafhoppers, Velvet Bean Caterpillars — Use 20 lb. per acre. Apply when larvae are young on Corn Earworms and Fall Armyworms. Apply when feeding by larvae is noted on Velvet Bean Caterpillars. Apply about July 10 and then at 3 week intervals on Potato Leafhoppers.

TOBACCO (PLANT BEDS): Tobacco Flea Beetles, Green June Beetle Larvae — Use 1-2 lb. per 100 sq. yds. Apply when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting.

TOBACCO FIELD: Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles — Use 20-40 lb. per acre, depending upon size of plants. Good coverage is essential. Use the highest dosage only on large plants. Repeat as needed. Allow 3 days before pruning or cutting.

COTTON: Early Season Control: Cotton Fleahopper, Cotton Leafworm, Flea Beetles, Thrips — Use 10 to 20 lb. per acre. **Mid and Late Season Control: Boll Weevils, Bollworms, Fall Armyworms, Leafhoppers, Tarnished Plant Bugs** — Use 20 to 40 lb. per acre, depending upon size of plants and insect population level. For Boll Weevil, apply every 5 to 7 days as required to maintain control.