

CAUTION

Keep Out of Reach of Children

WASH HANDS AND FACE


THOROUGHLY AFTER EACH APPLICATION

WASH HANDS AND FACE THOROUGHLY AFTER
EATING AND BEFORE EATING OR SMOKING

AVOID CONTAMINATION OF FEED OR FOODSTUFFS

BURN THE EMPTY CONTAINER COMPLETELY. EXERCISE
EXTREME CAUTION AND STAY WELL AWAY FROM THE SMOKE.

USDA Reg. No. 2299-175



NET WEIGHT — 50 POUNDS

ACTIVE INGREDIENTS:	
Pentachloronitrobenzene	10%
INERT INGREDIENTS	90%
Total	100%
TERRACLOR is a registered trademark of Chemical Corporation	

NOTICE Seller warrants that the product conforms to the chemical description and is reasonably fit for the uses stated on this label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

2-20-4-1

RECOMMENDATIONS

COTTON (Damping-Off, Seed and Seedling Rots — Rhizoctonia): FURROW APPLICATION — Apply 2 1/2 to 5 lb. per acre to the planting furrow. Dust over seed and soil at planting time, using suitable application equipment.

PLANTER BOX APPLICATION — COTTON SEED: Apply to the seed through the planter box at planting time at the rate of 6 to 10 lb. per acre. Thoroughly mix seed with seeds before planting. This is a supplementary treatment and should be used in addition to regular seed treatment.

Apply to the seed at planting rate in a rate planter for proper seeding and to avoid the suggested dosage of this product.

LAWNS (Brown Patch — Rhizoctonia Solani): For Southern or warm season lawns (e.g., Bermuda, Bahia, and similar grasses). Make one application in the fall, or spring, at first sign of infection. Apply 8 lb. per acre. After application, the treated area should be lightly watered to help movement through the grass to soil level. If the treated area is heavily trafficked, heavily mowed, or flooded, or if the disease is severe, reapplication may be necessary 3 to 4 weeks later.

SOUTHERN OR TROPICAL SEASON LAWNS SUCH AS RYE GRASS, FESCUE, BENT GRASS, AND SIMILAR GRASSES: Apply 2 1/2 to 5 lb. per 1,000 sq. ft. at first sign of infection, and repeat every 7 to 10 days during periods of hot weather. This product may be applied by spraying or watering, or by washing chemical down to the soil surface.

ROSES (Botrytis Storage Rot): Stack dormant roses in a dry, well-ventilated place. This causes rot and will disappear as the roses are planted.

Refer to recommendations on next page.

5905-310

10% TERRACLOR® DUST

PEANUTS (Southern Stem and Root Rot — Sclerotium Rolfii): PRE-PLANT SOIL MIX — Apply 100 to 150 lb. per acre in an 8 to 12 inch band centered on the row, and mix to a depth of 2 inches. Application may be made simultaneously with planting using suitable application equipment.

CAUTION: If a disease situation is severe, use very first sweeps of a treated band of Terraclor will remain active.

ORNAMENTALS

AFRICAN VIOLETS, BEGONIAS, CARNATIONS, CHRYSANTHEMUMS, POINSETTIAS, SNAPDRAGONS AND SIMILAR BEDDING PLANTS (Stem Rot — Rhizoctonia): Before Planting — Apply 15 to 20 lb. per 1,000 sq. ft. by dusting on the soil surface and thoroughly mixing into the top 2 inches of soil.

AZALEA (Petal Blight caused by Oculinia Azaleae): Apply 8 lb. per 1,000 sq. ft. as a ground dust. Begin applications prior to opening of buds and repeat every 3-4 weeks through bloom period.

period. Apply uniformly to the ground beneath the plants and surrounding area.

CAMELLIA (Flower Blight): To control this disease apply 2 1/2 to 5 lb. per acre to the surface of the soil or leaf litter beneath the plants. For smaller areas apply 5 lb. per 100 sq. ft. Begin application prior to opening of buds and repeat every 3 to 4 weeks through bloom period.

GLADIOLUS (Neck or Bulb Rot — Stromatinia or Sclerotinia Gladioli): BROADCAST — At planting time apply 1,200 lb. per acre (12 lb. per 100 sq. ft.) to soil surface and mix thoroughly to an upper 4 to 6 inches by double discing or rotovating. Bulbs may be planted immediately. ROW — At planting time apply 600 lb. per acre (6 lb. per 100 sq. ft.) in 8 to 10 inch band and thoroughly mix 4 to 6 inches deep by rotovating. Bulbs may be planted immediately.

ROSES (Botrytis Storage Rot): Stack dormant roses in a dry, well-ventilated place and apply by liberally dusting the branches immediately after stacking.

another premium product by

HELENA CHEMICAL COMPANY — SOUTHEAST

Tampa, Florida