

CAUTION

Keep Out of Reach of Children.

Harmful If Inhaled or Swallowed.

Do Not Breathe Dust. Do Not Take Internally.

Avoid Prolonged or Repeated Breathing of Dust.

Skin Contact May Be Harmful. Avoid Contact.

Wash Hands Thoroughly After Handling and Before Eating or Smoking. Take Shower or Bath After Work.

Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not contaminate food and feed.

Avoid drift to food and forage crops.


Consult State Agricultural Experiment Stations or State Agricultural Extension service for additional information, as timing, number and rate of applica-



10% SEVIN* 75% SULPHUR DUST

*Reg. TM of Union Carbide Corporation for the active ingredient, Carbaryl.

U.S. Patent No. 2,903,478



CAUTION KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTION STATEMENTS ON SIDE PANEL

ACTIVE INGREDIENTS:

Sulphur 75.00%
Carbaryl (1-naphthyl N-methylcarbamate) 10.00%

INERT INGREDIENTS 15.00%

TOTAL 100.00%

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

EPA Reg. No. 5905-311

USE ONLY AS RECOMMENDED

Completely empty container and bury unused chemicals at least 18 inches deep in an isolated location away from water supplies. Burn empty container completely. Exercise extreme caution. Stay away from smoke. Bury ashes, residue.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control. Observe use limitations given for specific crops.

RECOMMENDATIONS

Sevin Insecticide will not control insects if used with lime or other highly alkaline material.

PEANUTS: Armyworms, Alfalfa Caterpillars, Green Cloverworm, Leafhoppers, Thrips, Velvet Bean Caterpillar, Blister Beetles, Grasshoppers, Corn Earworm, Webworm, Stink Bugs; Leaf Spot - Use 10 to 15 pounds of dust per acre. Apply when insects first appear or Leaf Spot threatens. Repeat as required to maintain control.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any