

# 7.5% SEVIN - 75% SULPHUR DUST

## (INSECTICIDE-FUNGICIDE)

**ACCEPTED**  
FEB 13 1975  
Under the Federal Insecticide, Fungicide, and Rodenticide Act  
EPA Reg. No. 5905-312

**ACCEPTED**  
FEB 13 1975  
Under the Federal Insecticide, Fungicide, and Rodenticide Act  
EPA Reg. No. 5905-312

**CAUTION** KEEP OUT OF REACH OF CHILDREN.  
SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.



NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Active Ingredients	By Wt.
Carbaryl (1-methyl-2-naphthylcarbamate)	7.5%
Sulfur	75.0%
Inert Ingredients	17.5%

EPA Reg. No. 5905-312

\*For a Map of Under Control Cooperation for the active ingredient, for a list of Under Control agents.

U.S. Pat. No. 2,903,422



### CAUTION

Keep Out of Reach of Children

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

**CAUTION:** HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long-Sleeved Work Clothing. Change to Clean Clothing Daily.

Do not store or transport near feed or food. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHES AND WATERING RECEPTACLES. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local or State has proved that sulfur does not damage crops in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

KEEP PLUGS OF ALL CONTAINERS TIGHTLY CLOSED.  
BURN IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

### DIRECTIONS


This product is formulated for application by ground or air equipment.

### CAUTION

Sevin dust should not be used on insects if used with lime or other highly alkaline material.

BEANS: Mexican Bean Beetle, 10 lb. per acre. Leafhoppers, 15 lb. per acre. Green Bean, Powdery Mildew - 25 lb. per acre. Lygus Bugs (For California Only) - 28 lb. per acre. Apply evenly at first sign of infestation. When necessary reapply every 7 to 10 days. May be used up to harvest.

CORN: Corn Earworm, 10 lb. per acre. Apply at 3-day intervals. Do not apply to corn after tasseling. Use of this product in the tasseling period of corn causes the pollinated period to be shorter. Do not use on corn after tasseling. Make 4 to 5 applications at 7 to 10 day intervals.



(4)

COTTON: Early-Season—Thrips, Cotton Fleahopper--10 to 12 lbs. per acre as soon as insects appear. Mid and Late-Season -- Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator--20 to 25 lbs. per acre. Apply when insects appear and repeat at 5 to 7-day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

For concentrate dust, calibrate ground equipment or aerial dust apparatus accurately. After properly adjusting equipment to deliver the required low volume, maintain constant speed throughout application to insure even coverage. Strictly avoid excessive application.

To effectively apply concentrate dust by aircraft, use standard venturi-type equipment with spreader vanes or Swathmaster-type spreaders. Fly at normal height (4 to 6 feet above the top of the plants is suggested) and maintain regular swath width (usually 30 to 35 feet is indicated). For application with ground equipment, use tractor mounted power dusters with a metering device that feeds the concentrate dust through one nozzle per row (with deflector) or similarly designed electronic dusters.

GRAPES: Grape Leaf Elder--1st Brood: 15 to 20 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Broods: 20 to 25 lbs. per acre. Grape Leafhopper, Powdery Mildew-- 15 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Thorough coverage is essential. May be applied up to harvest.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Powdery Mildew--40 to 45 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Do not apply within 1 day of harvest.

PEANUTS: Amygdalus, Alabama Caterpillars, Green Cloverworms, Peanut Leafminers, Thrips, Velvet Bean Caterpillars, Blister Beetles, Grasshoppers, Corn Earworms, Weevils, Stink Bugs, Leaf Beetles-- Use 15 to 20 pounds of dust per acre. Apply when insects first appear or leaf spot threatens. Repeat as required to maintain control.

POPLARS: Colorado Potato Beetle--10 lbs. per acre. Leafhopper and Lygus Bugs-- 15 lbs. per acre. Lygus Bugs-- 25 lbs. per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10 day intervals.

TOMATOES: Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs--25 to 40 lbs. per acre. Russet Mite-- 40 lbs. per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the high content of this dust, first obtain permission from your carrier or processor before using it on tomatoes intended for sale to such carrier or processor.

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other expressed or implied warranty, including any other expressed or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages, such as loss of profits or value, etc.

