[5405-317]

## **CAUTION**

Keep Out of Reach of Children
HARMFUL IF INHALED OR SWALLOWED

Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust.


Skin Contact With Any Insecticide May Be Harmful.

Avoid Contact.

Wash Hands Thoroughly After Handling and Before Eating or Smoking. Take Shower or Bath After Work.

Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

Note for Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidatal.



**NET WEIGHT** 

50 POUNDS

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarba-

bamate) . . . . . . . . . . . . . . . 10.0% INTERT INGREDIENTS . . . . . . . . . . . . . . . 90.0%

TOTAL

100.0%

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably fore-seeable to seller, and buyer assumes the risk of any such

## OBSERVE FOLLOWING USE PRECAUTIONS

Sevin Insecticide will not control insects if used with lime or other highly alkaline materials

Taxic to honey bees. Do not apply during bloom when honey bees or other pollinators are active.

Avoid drift to adjoining food and forage crops.

The Federal Food and Drug Administration has set official tolerances for certain posticide residues in order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult state Agricultural Experiment Station for additional information, as the timing, number and rates of applications needed will vary with local conditions.

## 10% SEVIN\* DUST

\*Trade Mark of Union Carbido Corporation for the active ingredient carbaryl (1-naphthy). Neptethylombamate):

U.S. Patent No. 2,903 478

## RECOMMENDATIONS

Dosages given are in amounts of 10% CEVIN DUST per acre. Apply when insects first threaten damage to crop and repeat at 5 to 7 day intervals as required to maintain control.

COTTON: Boll Weevil, Bollworm, Cotton Fleahopper. Lygus Bugs, Pink Bollworm, Thrips. Early Season Control — Use 15 lb. per acre. Mid-Season — Use 12 to 20 lb. per acre. Late Season — Use 20 to 25 lb. per acre.

**PEANUTS:** Corn Earworm, Fall Armyworms, Potato Leafhoppers. Velvet Bean Caterpillars— Use 15 lb. per acre. Apply when larvae are young on Corn Earworm and Fall Armyworms. Apply when feeding by larvae is noted on Velvet Bean Caterpillars. Apply about July 10 and then at three-week intervals on leafhoppers.

TOBACCO (Plant Bed): Tobacco Flea Beetles, Green June Beetle Larvae — Uso 212 lb. per 100 sq. yds. Apply when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting.

TOBACCO (Field: Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles — Use 20 to 40 lb. per acre, depending upon the size of plants. Good coverage is essential. Use the highest dosage only on large plants. Repeat as needed. Allow 3 days before priming or cutting.

USE ONLY AS RECOMMENDED!

USDA Reg. No. 2299 209