

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED

Do Not Breathe Dust. Do Not Take Internally.

Avoid Prolonged or Repeated Breathing of Dust.

Skin Contact May Be Harmful. Avoid Contact.

Wash Hands Thoroughly After Handling and Before Eating or Smoking. Take Shower or Bath After Work.

Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control. Observe use limitations given for specific crops.


OBSERVE FOLLOWING USE PRECAUTIONS

Sevin Insecticide will not control insects if used with lime or other highly alkaline material.



10% SEVIN* - 3.4% COPPER
- 65% SULPHUR DUST

*Reg. TM of Union Carbide Corp.



CAUTION KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.

ACTIVE INGREDIENTS:

Copper, expressed as metallic.....	3.4%
Carbaryl (1-naphthyl N-methyl-carbamate).....	10.0%
Sulphur.....	65.0%
INERT INGREDIENTS:.....	21.6%
TOTAL.....	100.0%

The copper is in the form of copper oxide.

NET WEIGHT 50 POUNDS

EPA Reg. No. 5905-320

To protect honey bees, do not apply when crops are in bloom. Remove or cover hives if application is necessary during bloom. Avoid drift to food and forage crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified.

Consult State Agricultural Experiment Stations or State Agricultural Extension service for additional information, as timing, number and rate of applications needed will vary with local conditions.

RECOMMENDATIONS

PEANUTS: Potato Leafhopper, Thrips, Velvet Bean Caterpillar, Cercospora Leafspot - Apply at the rate of 10 pounds per acre when disease or insects first appear. If a heavy rain occurs within 24 hours of the first application repeat the application within three days. Repeat applications at 10 to 12 day intervals.

USE ONLY AS RECOMMENDED

Completely empty container and bury unused chemicals at least 18 inches deep in isolated location away from water supplies. Burn empty container completely. Exercise extreme caution. Stay away from smoke. Bury ashes, residue.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED

Do Not Breathe Dust. Do Not Take Internally.

Avoid Prolonged or Repeated Breathing of Dust.

Skin Contact May Be Harmful. Avoid Contact.

Wash Hands Thoroughly After Handling and Before Eating or Smoking. Take Shower or Bath After Work.

Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control. Observe use limitations given for specific crops.

OBSERVE FOLLOWING USE PRECAUTIONS


Sevin Insecticide will not control insects if used with lime or other highly alkaline material.

Do not contaminate any body of water or



10% SEVIN* - 3.4% COPPER
- 65% SULPHUR DUST

*Reg. TM of Union Carbide Corp.



CAUTION KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.

ACTIVE INGREDIENTS:

Copper, expressed as metallic.....	3.4%
Carbaryl (1-naphthyl N-methyl-carbamate).....	10.0%
Sulphur.....	65.0%
INERT INGREDIENTS:.....	21.6%
TOTAL.....	100.0%

The copper is in the form of copper oxide.

NET WEIGHT 50 POUNDS

EPA Reg. No. 5905-320

To protect honey bees, do not apply when crops are in bloom. Remove or cover hives if application is necessary during bloom. Avoid drift to food and forage crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified.

Consult State Agricultural Experiment Stations or State Agricultural Extension service for additional information, as timing, number and rate of applications needed will vary with local conditions.

RECOMMENDATIONS

PEANUTS: Potato Leafhopper, Thrips, Velvet Bean Caterpillar, Cercospora Leafspot - Apply at the rate of 10 pounds per acre when disease or insects first appear. If a heavy rain occurs within 24 hours of the first application repeat the application within three days. Repeat applications at 10 to 12 day intervals.

USE ONLY AS RECOMMENDED

Completely empty container and bury unused chemicals at least 18 inches deep in isolated location away from water supplies. Burn empty container completely. Exercise extreme caution. Stay away from smoke. Bury ashes, residue.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale.