

AREA

Extension Service or
ing treated areas.

STAINERS

pletely empty the contents
18 inches deep in an
supplies.
ainer and bury the pieces
ed location away from

fter into empty can and
detergent. Potate can
with the solution. Bury
p in an isolated location
les in top and bottom of
1 an isolated location.

ALLON DRUMS-Pour the
iners:

1/2 cup caustic soda
detergent.

iter, 1 lb. caustic soda

iter, 2 lb. caustic soda

urfaces are thoroughly
st 18 inches deep in an
plies.

ehold (ye) with extreme
on clothing. Read and
ackage. Flush eyes in
ontainer. Bury deeply in

CANNOT BE USED

burn in hot fire until all
off container. Extreme
re that containers are
s stay well away from
l to burn more than 5

AN

te in large doses. TWO
muscularly as soon as
to 10 minute intervals
2-PAM chloride is also
ed in conjunction with
IE OR TRANQUILIZERS.
e inhibitor affecting the
ystems and producing
in. At first sign of
ld be given supplemental
e. Continued absorption
al relapses have been
ent. VERY CLOSE
DICATED FOR AT LEAST

STOP-READ LABEL BEFORE USING!

POISON

See Side Panels
for Antidote &
Precautions



DANGER

Keep Out of
Reach of
Children



PARATHION 6 METHYL 3

(This product contains 6 lbs. Ethyl Parathion &
3 lbs. Methyl Parathion per gallon.)

ACTIVE INGREDIENTS:

O,O-dimethyl O-p-nitrophenyl phosphorothioate*	29.03%
Parathion (O,O-diethyl O-p-nitrophenyl phosphorothioate)	57.77%

INERT INGREDIENTS 13.20%

TOTAL 100.00%

*Methyl Parathion

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE

MANUFACTURED BY

RECOMMENDATIONS

VEGETABLES

To insure that residue levels are below tolerances established by the U.S. Food & Drug Administration, do not apply to any crop at a rate higher than specified.

DO NOT APPLY WITHIN 5 DAYS OF HARVEST ON THE FOLLOWING: **EQUISETACEAE:** For control of aphids, blister beetles, leafhoppers, leaf miners, plant bugs, potato phylloxera, thrips and vegetable weevils, use 1/2 to 1 1/3 pts. per acre. **DO NOT APPLY WITHIN 12 DAYS OF HARVEST ON FOLLOWING:**

CORN: (Seed, Corn, Popcorn): For control of rootworm adults, use 1/3 pt. per acre. For aphids, armyworms and fall armyworms up to 3rd instar, budworms, spider mites, climbing cutworms, and European corn borers, use 1/2 pt. per acre. For stink bugs, use 2/3 pt. per acre. For European corn borer control, apply first spray when 75% of corn plants show SHOT HOLE feeding and follow with 2 additional sprays at 5 to 7 day intervals. Use sufficient water to provide complete coverage and be certain whorls of plants are well treated. **DO NOT APPLY WITHIN 15 DAYS OF HARVEST ON FOLLOWING:** (When tops of Beets and Turnips are used as food, do not apply within 21 days of harvest.)

BEETS: (Leaf) For control of aphids, blister beetles and webworms, use 1/2 pt. per acre. Do not apply within 21 days of harvest. **CARROTS:** For control of aphids, vegetable weevils, leaf hoppers and potato beetles, use 1/2 pt. per acre. Do not use carrot tops for food or feed.

CUCUMBERS: For control of aphids and two-spotted mites, use 1/2 pt. per acre. Do not apply before vining. **PEAS:** (Garden and Blackeye): For control of aphids and potato beetles, use 1/2 pt. per acre. For armyworms up to 3rd instar, leafhoppers and spider mites, use 2/3 pts. per acre. **PEPPERS:** For control of aphids and leaf miner, use 1/2 pt. per acre. For flea beetles, use 2/3 to 3/4 pts. per acre. **DO NOT APPLY WITHIN 21 DAYS OF HARVEST ON FOLLOWING:**

COLLARD'S KALE, MUSTARD: For control of aphids, Cabbage looper, imported leaf miner, vegetable weevils, leafhopper, and cucumber beetles, use 1/2 pt. per acre. **SPINACH:** For control of aphids, cabbage looper, imported cabbage worms, leaf miners, vegetable weevils, leafhopper, crow flies and sand corn maggot, use 2/3 pt. per acre. **LETTUCE (Leaf, Bibb Head):** For control of aphids, armyworms up to 3rd instar, imported cabbage worms, use 1/2 pt. per acre. For cabbage looper, use 2/3 pt. per acre. **DO NOT APPLY WITHIN 30 DAYS OF HARVEST.**

CELERY: For control of aphids, celery leaf tiers, celery worm, spider mites and tarnished plant bugs, use 1/2 pt. per acre. For leafhoppers, use 3/4 pt. per acre.

FIELD AND FORAGE CROPS

COTTON: For control of cotton leafworms, leafhoppers, and spider mites, thrips and garden webworms, use 1/2 pt. per acre.

For control of cabbage looper, bollworms and budworms, aphids, cutworm leaf perforators, armyworms up to 3rd instar, stink bugs, Lygus bugs and saltmarsh caterpillars, use 1 pt. per acre. Application to cotton should be made at 4 to 5 day intervals until control is obtained. Caution: When necessary to enter treated fields within 48 hours after application, protective clothing should be worn. Do not apply within 7 days of harvest or hand-picking cotton.

CORN: (Hybrid Seed, Corn, Field Corn): For control of rootworm adults, use 1/3 pt. per acre. For aphids, armyworms and fall armyworms up to 3rd instar, budworms, spider mites, climbing cutworms and European corn borers, use 1/2 pt. per acre. For stink bugs, use 2/3 pt. per acre. For European corn borer control, apply first spray when 75% of corn plants show SHOT HOLE feeding and follow with 2 additional sprays at 5 to 7 day intervals. Use sufficient water per acre to provide complete coverage and be certain whorls of plants are well treated. Do not apply within 12 days of harvesting, cutting for forage use. **SUGAR BEETS:** For control of alfalfaoppers, aphids, armyworms up to 3rd instar, blister beetles, flea beetles, leafhoppers, leaf miners, Lygus bugs, stink bugs and webworms, use 1/2 to 2/3 pt. per acre. If tops are to be used for cattle feed, do not apply within 20 days of harvest. Do not apply within 60 days of harvest for forage use. **SORGHUM:** For control of corn earworms, corn leaf aphids, large chinch bugs and sorghum webworms, use 1/3 pt. per acre.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear water proof pants, coat, hat, rubber boots, rubber overshoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the U.S. Bureau of Mines for Parathion protection. (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK, take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

POISON SIGNS (Symptoms)

Parathion is every dangerous poison. It rapidly enters the body by contact with all skin surfaces and eyes. Clothing in contact with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling, frothing of mouth and nose, muscle spasms and coma.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risk referred to therein.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except writing with a specific reference to this warranty. All damages arising from a breach of this warranty shall be limited to direct damages, and not include consequential or commerial damages such as loss of profit or value, etc.

NOT FOR HOME USE